

Stable non-BPS states in string theory: a pedagogical review ^{*}

Alberto Lerda ^a and Rodolfo Russo ^b

^a *Dipartimento di Scienze e Tecnologie Avanzate
Università del Piemonte Orientale (Alessandria)*

*and Dipartimento di Fisica Teorica, Università di Torino
and I.N.F.N., Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy*
lerda@to.infn.it

^b *Institut de Physique, Université de Neuchâtel
Rue A.-L. Breguet 1, CH-2000 Neuchâtel, Switzerland*
Rodolfo.Russo@iph.unine.ch

Abstract

We present a pedagogical review of the stable non-BPS states in string theory which have recently attracted some attention in the literature. In particular, following the analysis of A. Sen, we discuss in detail the case of the stable non-BPS D-particle of Type I theory whose existence is predicted (and required) by the heterotic/Type I duality. We show that this D-particle originates from an unstable bound state formed by a D1/anti-D1 pair of Type IIB in which the tachyon field acquires a solitonic kink configuration. The mechanism of tachyon condensation is discussed first at a qualitative level and then with an exact conformal field theory analysis.

^{*}This work is partially supported by the European Commission TMR programme ERBFMRX-CT96-0045 and by MURST, and is based on the lectures given at the TMR school *Quantum Aspects of Gauge Theories, Supersymmetry and Unification* in Leuven (Belgium) January 18-23 1999.

1 Introduction

During the last few years our understanding of string theory has dramatically increased. Nowadays it is known that all five consistent superstring models (*i.e.* Type IIA, Type IIB, Type I, $SO(32)$ heterotic and $E_8 \times E_8$ heterotic [1]) and the eleven dimensional supergravity are different perturbative expansions of a unique underlying theory, called M-theory. However, despite numerous attempts and remarkable achievements, we are still far from a complete and satisfactory formulation of M-theory. The best we can do at present time is to define it by means of the non-perturbative duality relations that connect the various corners of its moduli space corresponding to the different string theories. Establishing (and checking) these relations precisely is therefore a necessary step towards a better understanding of M-theory.

One way of testing a duality relation between theory A and theory B is to show that their effective actions are equivalent up to a field redefinition. In this way, for example, it was discovered that Type IIB superstring is self-dual [2], or that the $SO(32)$ heterotic string is dual to Type I theory [3]. Of course, by focusing on the effective actions, one checks a duality only at a field theory level, and not in a really stringy way. However, a deeper understanding can be obtained by looking at the so-called BPS states. These are a special class of states, which are characterized by the key property that their mass is completely determined by their charge under some gauge field. They form short (or ultra-short) multiplets of the supersymmetry algebra of the theory, and because of this fact they are *stable* and protected from quantum radiative corrections. Therefore, their properties can be studied and analyzed perturbatively at weak coupling in the theory A, and then safely extrapolated at strong coupling where they can be reinterpreted in terms of non-perturbative configurations of the dual theory B. A well-known example of such BPS states is given by the supersymmetric Dp branes of Type II theories (with p even in Type IIA and p odd in Type IIB) which are charged under the $(p+1)$ gauge potentials of the R-R sector [4, 5]. Since the BPS configurations are protected by non-renormalization theorems, one may legitimately pose the question whether the non-perturbative tests based on them are really a strong evidence for a duality conjecture or simply unavoidable results dictated by supersymmetry alone. Thus, it is of great conceptual importance to analyze a non-perturbative duality beyond the BPS level.

To this aim, it is useful to observe that quite often a string theory contains in its spectrum states which are *stable* without being BPS. These are in general the lightest states which carry some conserved quantum numbers of the theory. For these states there is no particular relation between their mass and their charge; they receive quantum corrections and form long multiplets of the supersymmetry algebra of the theory (*i.e.* they are non-BPS). However, being the lightest states with a given set of conserved quantum numbers, they are stable since they cannot decay into anything else. In general it is not difficult to find such stable states with the standard perturbative methods of string theory and analyze their properties at weak coupling; but then a very interesting question can be asked: what happens to these stable states at strong coupling? Since they cannot decay, these states should be present also in the strong coupling regime, and thus, if the string duality relations are correct, one should find stable non-BPS states with the right multiplicities and quantum numbers also in the dual theory. To verify the existence of such states is therefore a very strong (perhaps the strongest) check on a non-perturbative duality relation between two string theories.

In these notes, following the analysis of A. Sen [6, 7], we apply this idea to test the heterotic/Type I duality [3] at a non-BPS level ¹. In this way we get a more complete understanding of this non-perturbative relation which, in some sense, has a privileged status: indeed, a study of the web of string dualities [12] shows that all of them can be “derived” from it by combining it with T-duality.

At the first massive level the $SO(32)$ heterotic string contains perturbative states which are stable but not BPS. Their stability follows from the fact that they are the lightest states carrying the quantum numbers of the spinor representation of the gauge group. Since they cannot decay, these states should be present also in the strong coupling regime. Then, if the heterotic/Type I duality is correct, Type I theory should support non-perturbative stable configurations that are spinors of $SO(32)$. It turns out [13] that a pair formed by a D1-brane and an anti D1-brane of Type I (wrapped on a circle and with a Z_2 Wilson line) describes a configuration with the quantum numbers of the spinor representation of $SO(32)$. Thus this system is the right candidate to describe in the non-perturbative regime the stable non-

¹A first check of this duality has been given in Ref. [8] where it has been shown that the world-volume theory on the D-string of Type I is identical to the world-sheet theory of the heterotic string. Later, further tests involving BPS saturated terms have been performed in Ref. [9] for the uncompactified case and, for the compactified case, in Ref. [10] at level of BPS spectra and in Ref. [11] at level of the effective actions.

BPS states of the heterotic string mentioned above. However, a superposition of a brane with an anti-brane is unstable due to the presence of tachyons in the open strings stretching between the brane and the anti-brane [14]. Then the problem of defining properly this superposition and treating its tachyon instability arises. This has been addressed by A. Sen in a remarkable series of papers [15, 16, 17, 6, 7, 18]. In particular in Ref. [6] he considered a D-string/anti-D-string pair of Type IIB, and managed to prove that when the tachyon condenses to a kink along the (compact) direction of the D-string, the pair becomes tightly bound and, as a whole, behaves as a D-particle. He also computed its mass finding that it is a factor of $\sqrt{2}$ bigger than the one of the supersymmetric BPS D-particle of Type IIA theory. The presence of a D-particle in Type IIB spectrum looks surprising at first sight since one usually thinks that in Type IIB there are only Dp branes with p odd. However, one should keep in mind that such a D-particle is a non-supersymmetric and non-BPS configuration. Furthermore it is unstable, due to the fact that there are tachyons in the spectrum of the open strings living on its world-line. These tachyons turn out to be odd under the world-sheet parity Ω , and hence disappear if one performs the Ω projection to get the Type I string [19]. Therefore, the D-particle found by Sen is a stable non-perturbative configuration of Type I that transforms as a spinor of $SO(32)$.

Another interesting example of stable non-BPS configurations that can be used to check a non-perturbative duality relation is provided by the states that live on the world-volume of a D5 brane of Type IIB on top of an orientifold $O5$ plane [13, 16, 20]. The theory which supports these states is obtained by modding out Type IIB theory in ten dimensions by $\Omega\mathcal{I}_4$ where Ω is the world-sheet parity (the twist operator) and $\mathcal{I}_4$ is the space-time parity which changes the sign of four of the space-like coordinates, *i.e.*

$$\text{IIB}/\Omega\mathcal{I}_4 . \tag{1.1}$$

This theory contains 32 D5-branes which are pairwise located on top of 16 orientifold 5-planes. The open strings that start and end on the two mirror D5 branes of a given orientifold plane are charged under an abelian gauge potential. It turns out that the lightest states which carry this abelian charge are at mass-level one, and thus are non-BPS but stable. To describe these states at strong coupling, we perform an S-duality transformation which maps the twist parity Ω to $(-1)^{F_L}$ where F_L is the left part of the space-time fermion number operator. Thus, the S-dual theory of (1.1) is the orbifold

$$\text{IIB}/(-1)^{F_L}\mathcal{I}_4 , \tag{1.2}$$

which contains the usual untwisted sector together with a twisted sector localized on the orbifold fixed plane. In particular at the massless level there are, among other fields, a 2-form Ramond-Ramond potential $A^{(2)}$ arising from the untwisted R-R sector and a 1-form Ramond-Ramond potential $A^{(1)}$ arising from the twisted R-R sector. It turns out (see Ref. [16] for details) that the consistent BPS states of this orbifold theory must be charged both under $A^{(2)}$ and under $A^{(1)}$. Therefore, they are D1-branes which carry also the charge of an abelian vector potential. Given this fact, it is then natural to conjecture that the non-BPS states which we are looking for can be obtained as a superposition of one D1-brane and one anti-D1-brane of this kind. Indeed, by taking this combination we cancel the charge of $A^{(2)}$ and obtain a solitonic configuration which carries only the charge of an abelian vector potential and behaves as a D-particle. This construction has been performed in Refs. [16, 20] using the boundary state formalism [21] in which the D-branes are described by a closed string state (the boundary state) that inserts a boundary on the string world-sheet and enforces on it the appropriate boundary conditions ². In particular, after treating properly the tachyon instability on the brane/anti-brane pair in the orbifold model (1.2), a stable D-particle state has been explicitly constructed and its mass has been computed. Of course a similar analysis can be done also in other orbifold models [23, 15] which involve D-branes of different dimensionality (see for a review Ref. [24]).

Like in the heterotic/Type I case, also in the orbifold theory (1.2) one finds that the stable non-BPS states predicted by duality arise from an unstable system formed by a D-string/anti-D-string pair of Type IIB in which the tachyon field acquires a solitonic kink configuration. Thus, one may wonder whether this mechanism is peculiar to the two cases we have mentioned, or can occur in more general situations. This question has been considered recently by several authors [25, 26, 18] using different methods, and new kinds of relation between BPS and non-BPS branes have been discovered. It is natural to hope that they may provide new ideas and clues to understand better the various D-branes and their fundamental role in string (or M) theory.

For Type II theories these new relations are represented in Fig. 1. Let us start from a pair formed by a Dp brane and an anti- Dp brane of Type IIB (*i.e.* p odd) which, as mentioned above, is an unstable system due to the presence of an open-string tachyon on its world-volume. If we condense this tachyon into a solitonic kink

²For a general discussion of the boundary state formalism and its application to the study of D-branes and their interactions see for example Ref. [22].

solution, then the resulting bound state describes a non-BPS $D(p-1)$ brane of Type IIB [6, 25, 18]. In these notes we will discuss in some detail only the case $p = 1$ which is relevant of the heterotic/Type I duality, but the same arguments can be used in all other cases as well. The non-BPS $D(p-1)$ brane of Type IIB constructed in this way is unstable because its world volume theory contains again a tachyonic excitation (which has nothing to do with the previous one). If we repeat the procedure and make this second tachyon field condense into a solitonic kink solution, we obtain as a result a $D(p-2)$ brane of Type IIB, which is supersymmetric and stable [18]. These descent relations, which of course can be established also with the D-branes of Type IIA theory, are represented by the vertical arrows in Fig. 1.




Figure 1: This figure resumes the relations between different D-branes in type-II superstring theories. The squares represent the usual supersymmetric BPS D-branes, while the circles stay for the non-BPS configurations (unstable in Type II theories). Starting from a pair formed by a D_p and an anti- D_p brane, a non-BPS brane can be constructed in two ways: one can mod out the system by $(-1)^{F_L}$ (horizontal arrows) or condense the tachyon living on its world-volume (vertical arrows). By repeating these operations twice, one finds a supersymmetric configuration. The diagonal links represent the usual T-duality.

Now let us consider another kind of relations between BPS and non-BPS branes. Again we start from the Dp /anti- Dp pair of Type IIB and see what happens when we mod out the system by $(-1)^{F_L}$ ³. A detailed analysis of the open strings that live on its world-volume shows that the configuration one obtains after modding out $(-1)^{F_L}$ has the same features of a Dp brane of Type IIA. Since p is odd, this must be a non-supersymmetric and non-BPS brane. By modding out once more by $(-1)^{F_L}$ one goes back to Type IIB theory and finds a stable and supersymmetric Dp brane [18, 27]. These relations are represented by the horizontal arrows in Fig.1. By exploiting all these relations (as well the usual T-duality) we can conclude that actually all branes of Type II theories descend from a bound state of D9 and anti-D9 branes, a fact that has been used in Refs. [25, 26] to show that the D-brane charges take values in the K-theory of space time (see also Refs. [28, 29, 30]).

These notes are organized as follows: in Section 2 we briefly summarize the essential perturbative features of the $SO(32)$ heterotic string and of the Type I theory; in Section 3 we study in some detail the D1 brane of Type I and the open string excitations that live on its world-sheet and discuss their relations with the heterotic theory. In Section 4 we show how a stable D-particle configuration of Type I can be constructed from an unstable D1/anti-D1 pair and study the mechanism of tachyon condensation from the effective field theory point of view. In Sections 5 and 6 we put these results on more solid ground by providing an exact conformal description of the non-BPS D-particle, and finally in Appendix A we collect a few technical details on the bosonization procedure which is used in the conformal field theory analysis. Most of our presentation in the last sections is based in an essential way on the series of papers by A. Sen [15, 16, 17, 6, 7, 18].

³The operator $(-1)^{F_L}$ acts on the closed string states by simply changing the sign of every state that has a left-moving part in the R sector. Thus, it is an exact symmetry of Type II theories; in particular $(-1)^{F_L}$ transforms a Dp brane into an anti- Dp brane and viceversa. Therefore a brane/anti-brane pair is invariant under this operator and so it is meaningful to ask what happens when we mod it out.

2 The $SO(32)$ heterotic string and the Type I theory: a brief perturbative description

In this section we are going to summarize the essential perturbative features of the $SO(32)$ heterotic string and of the Type I theory that will be relevant for our later discussions. In subsection 2.1 we briefly analyze the spectrum of the heterotic string and show that the states in the spinor representation of $SO(32)$ are non-BPS but stable. Then, in subsection 2.2 we describe the content of the Type I string viewed as an orientifold of Type IIB theory. In particular we focus on the open string sector and also discuss the main properties of the so-called 1-1 strings that live on the world-sheet of the D1 branes of the theory.

2.1 The $SO(32)$ heterotic string

The heterotic string is a theory of closed oriented strings with a local gauge invariance in ten dimensions. Roughly speaking, it can be considered as a combination of the bosonic string and the superstring. The basic idea of the heterotic construction is to exploit the holomorphic factorization between the left and the right moving modes of closed strings and to treat them in two different ways. In particular, in the right sector one introduces the usual field content of the superstring, *i.e.* bosonic and fermionic coordinates $\widetilde{X}^\mu(\tau-\sigma)$ and $\widetilde{\psi}^\mu(\tau-\sigma)$ with $\mu = 0, 1, \dots, 9$ (plus the corresponding ghosts and superghosts), and performs the standard GSO projection. In the left sector, instead, one introduces only bosonic coordinates $X^\rho(\tau+\sigma)$ with $\rho = 0, 1, \dots, 25$ (plus the corresponding ghosts). Since not all X^ρ 's can be interpreted as spacetime coordinates, the left-moving sector is naturally divided in two sets: when $\rho = 0, \dots, 9$, the fields $X^\rho(\tau+\sigma)$ are combined with $\widetilde{X}^\mu(\tau-\sigma)$ to build the usual closed string coordinates $X^\mu(\tau, \sigma)$, while for $\rho = 10, \dots, 25$, the left-moving fields $X^\rho(\tau+\sigma)$ are interpreted as internal degrees of freedom that can propagate on the world-sheet. We will label these sixteen internal coordinates with an index $A = 1, \dots, 16$.

Given this field content, it is easy to realize that a typical element of the Fock space of the heterotic theory is a tensor product of a left and a right state, and looks like

$$\left(\alpha_{-n_1}^{\mu_1} \cdots \alpha_{-n_\ell}^{A_\ell} \cdots |k; p\rangle\right) \otimes \left(\widetilde{\alpha}_{-m_1}^{\nu_1} \cdots \widetilde{\psi}_{-r_p}^{\nu_p} \cdots |\widetilde{k}\rangle\right) \quad (2.1)$$

where, in an obvious notation, we have denoted by α_{-n} and $\tilde{\alpha}_n$ the creation modes of the bosonic fields and by $\tilde{\psi}_{-r}$ the creation modes of the fermionic coordinates. These ones have half-integer indices in the NS sector and integer indices in the R sector. Notice that the left vacuum in Eq. (2.1) depends on the spacetime momentum k^μ as well as on the internal momentum p^A , whereas the right vacuum depends only on k^μ . If one considers states in the R sector, the right vacuum will carry also a spinor label; in these notes however, we will not consider explicitly this case and focus only on the NS sector.

The states of the form (2.1) are not all acceptable in the heterotic theory; indeed a physical state $|v\rangle$ must satisfy the Virasoro constraints. In particular the mass-shell condition is

$$(L_0 - 1)|v\rangle = \left(\tilde{L}_0 - \frac{1}{2}\right)|v\rangle = 0 \quad , \quad (2.2)$$

where we have used the values of the intercept of the bosonic theory ($a = -1$) in the left sector and of the NS superstring ($a = -1/2$) in the right sector. By expanding $\tilde{L}_0$ in modes, one easily finds from the second equality in Eq. (2.2) that the mass M of a state is given by

$$M^2 = \frac{4}{\alpha'}\left(\tilde{N} - \frac{1}{2}\right) \quad (2.3)$$

where $\tilde{N}$ is the total number of right moving oscillators. From the first equality of Eq. (2.2) one can derive a generalized level matching condition which relates $\tilde{N}$ to the total number of left moving oscillators N that are present in a given state. This condition reads as follows

$$\tilde{N} + \frac{1}{2} = N + \frac{1}{2} \sum_A (p^A)^2 \quad , \quad (2.4)$$

where we have conveniently measured the internal momenta p^A in units of $\sqrt{2\alpha'}$. Further restrictions come from the GSO projection that one has to perform in the right sector to have a consistent model. In particular, the GSO projection on NS states selects only half-integer occupation numbers $\tilde{N}$. Since the left occupation number N is always integer as in the bosonic theory, in order to be able to satisfy Eq. (2.4), the internal momenta p^A have to be quantized. In particular, the quantity $\sum_A (p^A)^2$ has to be an even number. This condition implies that the internal coordinates X^A must be compactified on an *even* 16-dimensional lattice. The intrinsic consistency of the theory, and more specifically the modular invariance of the one-loop partition function, requires that this lattice be also *self-dual*. It can be shown that there exist only two 16-dimensional lattices satisfying both these

properties: the root lattice of $E_8 \times E_8$, and a Z_2 sublattice of the weight lattice of $SO(32)$. Since we are interested in the heterotic theory with gauge group $SO(32)$, for the rest of these notes, we will focus only on the second lattice which is denoted by Γ_{16} and is defined by

$$(n_1, \dots, n_{16}) \in \Gamma_{16} \quad \text{and} \quad \left(n_1 + \frac{1}{2}, \dots, n_{16} + \frac{1}{2}\right) \in \Gamma_{16} \quad \Longleftrightarrow \quad \sum_i n_i \in 2Z \quad (2.5)$$

It is interesting to observe that even if in the original formulation only a $SO(16)$ symmetry is manifest, when the internal coordinates are compactified on Γ_{16} , the resulting gauge group is much bigger. This fact can be explicitly checked by counting how many massless states involving the internal coordinates X^A are present in the theory. Because of Eq. (2.3), all such states must have $\tilde{N} = 1/2$, so that their right-moving part is simply $\tilde{\psi}_{-1/2}^i |\tilde{k}\rangle$ where $i = 2, \dots, 9$ labels the directions transverse to the light-cone. On the other hand, the level matching condition (2.4) requires that

$$N + \frac{1}{2} \sum_A (p^A)^2 = 1 \quad (2.6)$$

This condition can be satisfied either by taking $N = 1$ and $p^A = 0$, or by taking $N = 0$ and the momenta p^A to be of the form $P = (\pm 1, \pm 1, 0, \dots, 0)$ (or any permutation thereof). With the first choice, we construct 16 states $\alpha_{-1}^A |k; 0\rangle$, while with the second choice we construct 480 states $|k; P\rangle$. Altogether we have 496 massless states that carry a spacetime vector index from the right-moving part and span the adjoint representation **496** of $SO(32)$. Thus, they can be legitimately identified with the gauge fields of $SO(32)$. The fact that a system of bosonic fields compactified on an appropriate space exhibits an enlarged symmetry can be understood also from a different point of view through the bosonization procedure [31] (see also Appendix A for some details). In fact by exploiting the equivalence between a compact boson and two fermions, one can substitute the original coordinates X^A with 32 fermions Λ^I ($I = 1, \dots, 32$) directly in the sigma model lagrangian of the heterotic theory, thus making manifest the $SO(32)$ symmetry.

It is important to realize that there are 64 more bosonic massless states in the theory which correspond to $\tilde{N} = 1/2$, $N = 1$ and $p^A = 0$, and are given by

$$\alpha_{-1}^i |k; 0\rangle \otimes \tilde{\psi}_{-1/2}^j |\tilde{k}\rangle \quad (2.7)$$

where the indices i, j run along the transverse directions $2, \dots, 9$. These states are clearly singlets with respect to the gauge group but are space-time tensors.

Decomposing them into irreducible components, we get a graviton, a dilaton and an antisymmetric two-index tensor. In conclusion, the bosonic massless states of the heterotic theory are in the following representations

$$(\mathbf{1}; \mathbf{1}) \oplus (\mathbf{35}; \mathbf{1}) \oplus (\mathbf{28}; \mathbf{1}) \oplus (\mathbf{8}; \mathbf{496}) \quad (2.8)$$

where in each term the two labels refer to the Lorentz and gauge groups respectively. By analyzing the R sector, one finds an equal number of fermionic massless states that complete the $N = 1$ supersymmetric multiplets.

Let us now consider the first excited level of the NS sector that consists of states with $\widetilde{N} = 3/2$ and mass squared $M^2 = 4/\alpha'$. There are 128 ways to realize $\widetilde{N} = 3/2$, namely

$$\widetilde{\psi}_{-3/2}^i |\widetilde{k}\rangle \rightarrow 8 \text{ states} \quad (2.9)$$

$$\widetilde{\alpha}_{-1}^i \widetilde{\psi}_{-1/2}^j |\widetilde{k}\rangle \rightarrow 64 \text{ states} \quad (2.10)$$

$$\widetilde{\psi}_{-1/2}^i \widetilde{\psi}_{-1/2}^j \widetilde{\psi}_{-1/2}^\ell |\widetilde{k}\rangle \rightarrow 56 \text{ states} \quad (2.11)$$

These massive states represent a symmetric two-index tensor and an antisymmetric three-form under the Lorentz group. In fact, the antisymmetric part of Eq. (2.10) together with the states (2.11) completes a massive three-form transforming in the representation **84** of the Lorentz group, while the remaining states transform together as a symmetric two-index tensor in the representation **44**. The level matching condition (2.4) requires to make a tensor product with left-moving configurations such that

$$N + \frac{1}{2} \sum_A (p^A)^2 = 2 \quad (2.12)$$

There are 73,764 ways to satisfy this requirement! The complete list of the corresponding states can be found for example on pag. 342 of Ref. [1] where it is shown that they transform as scalars, spinors, second-rank antisymmetric tensors, fourth-rank antisymmetric tensors and second-rank symmetric traceless tensors of $SO(32)$. Here we focus on the 2^{15} states that are obtained by taking in Eq. (2.12) $N = 0$ and momenta p^A of the form $(\pm\frac{1}{2}, \pm\frac{1}{2}, \dots, \pm\frac{1}{2})$ with an even number of plus signs. Notice that these momenta define a point in the lattice Γ_{16} , since they satisfy the second condition of Eq. (2.5), and correspond to the *spinor* representation of $SO(32)$. Indeed, similarly to what happens with the usual spin fields, half-integer valued momenta always produce states which are spinors of the symmetry group. By combining these left modes with the right-moving ones of Eqs. (2.9) - (2.11),

we then obtain bosonic states transforming as

$$(44; \mathbf{2}^{15}) \oplus (84; \mathbf{2}^{15}) \quad . \quad (2.13)$$

Analyzing the first excited level of the R sector, one can find 128 massive fermionic states which transform in the spinor representation of $SO(32)$ and complete the $N = 1$ supersymmetry multiplets. Thus, altogether the spinors of $SO(32)$ appear with 256 different polarizations, 128 bosonic and 128 fermionic, which correspond to a *long* multiplet of the $N = 1$ supersymmetry algebra in ten dimensions. Thus, these states cannot satisfy the BPS condition [32] but are, nevertheless, stable. Indeed, as we have seen, at the massless level there are no spinors of $SO(32)$ and thus the states we have found are the lightest ones with these quantum numbers and hence cannot decay. As it was observed on pag. 343 of Ref. [1], this feature makes these spinors “absolutely stable, which might be an interesting prediction for the $SO(32)$ heterotic theory”. This expectation is going to be confirmed within the non-perturbative duality between the $SO(32)$ heterotic string and the Type I theory.

2.2 The Type I string

The Type I string theory is a $N = 1$ supersymmetric model in ten dimensions that consists of both open and closed *unoriented* strings with the usual GSO projection⁴. It is by now well-known that the Type I theory can be interpreted as an orientifold of Type IIB theory, which is a chiral model consisting only of closed oriented strings [19]. For a general discussion on the orientifold projection we refer for example to Ref. [13]. Here it is sufficient to recall that the orientifold projection is defined by means of a discrete transformation Ω which in general can mix the world-sheet parity with a space-time operation. Then, a new consistent spectrum is obtained by removing all states of the original theory that are not invariant under Ω , and by adding new states that are defined by a generalized periodicity condition on the string coordinates (the so-called twisted sector).

For the Type I theory, the operator Ω is simply the world-sheet parity, namely

$$\Omega \quad : \quad \sigma \longrightarrow \pi - \sigma \quad (2.14)$$

⁴Note that for closed unoriented strings the GSO projector has to be chiral, like the one of Type IIB string.

From this definition, it is easy to derive the action of Ω on the string modes; in fact, by expanding the closed string coordinates in the usual manner

$$X^\mu(\tau, \sigma) = x^\mu + 2\alpha' p^\mu \tau + i\sqrt{\frac{\alpha'}{2}} \sum_{n \neq 0} \left[\frac{\alpha_n^\mu}{n} e^{-2in(\tau+\sigma)} + \frac{\tilde{\alpha}_n^\mu}{n} e^{-2in(\tau-\sigma)} \right] , \quad (2.15)$$

one immediately sees that Ω interchanges the left and the right moving oscillators, *i.e.*

$$\Omega \alpha_n^\mu \Omega^{-1} = \tilde{\alpha}_n^\mu \quad , \quad \Omega \tilde{\alpha}_n^\mu \Omega^{-1} = \alpha_n^\mu \quad (2.16)$$

World-sheet supersymmetry requires that also the fermionic coordinates $\psi^\mu(\tau + \sigma)$ and $\tilde{\psi}^\mu(\tau - \sigma)$ transform in a similar way, *i.e.*

$$\Omega \psi_r^\mu \Omega^{-1} = \tilde{\psi}_r^\mu \quad , \quad \Omega \tilde{\psi}_r^\mu \Omega^{-1} = \psi_r^\mu \quad (2.17)$$

where the moding index r is half-integer in the NS sector and integer in the R sector. To proceed further, we have to specify also the action of Ω on the ground states. For simplicity, we choose the $(-1, -1)$ superghost picture in the NS-NS sector and the $(-1/2, -1/2)$ superghost picture in the R-R sector [31]. Since on the matter fields Ω simply exchanges the left and the right sector, it is natural to assume that

$$\Omega \left(|0\rangle_{-1} \otimes |\tilde{0}\rangle_{-1} \right) = |\tilde{0}\rangle_{-1} \otimes |0\rangle_{-1} = -|0\rangle_{-1} \otimes |\tilde{0}\rangle_{-1} \quad (2.18)$$

for the NS-NS sector, and

$$\Omega \left(|S^\alpha\rangle_{-1/2} \otimes |\tilde{S}^\beta\rangle_{-1/2} \right) = |\tilde{S}^\alpha\rangle_{-1/2} \otimes |S^\beta\rangle_{-1/2} = -|S^\beta\rangle_{-1/2} \otimes |\tilde{S}^\alpha\rangle_{-1/2} \quad (2.19)$$

for the R-R sector, with S^α and $\tilde{S}^\beta$ being the left and right spin fields in the chiral spinor representation of $SO(1, 9)$. Notice that the $-$ signs in Eqs. (2.18) and (2.19) originate from the exchange of the left and right vacua which have fermionic statistics.

Using these rules, it is easy to see which states of Type IIB theory are even under Ω and must be kept, and which states are odd and must be removed. Let us consider for example the massless states of the NS-NS sector which are represented by

$$|\Phi\rangle = \epsilon_{\mu\nu} \psi_{-1/2}^\mu \tilde{\psi}_{-1/2}^\nu |0\rangle_{-1} \otimes |\tilde{0}\rangle_{-1} \quad (2.20)$$

where the polarization tensor satisfies $k^\mu \epsilon_{\mu\nu} = \epsilon_{\mu\nu} k^\nu = 0$ with $k^2 = 0$. Then, using Eqs. (2.17) and (2.18) we can obtain

$$\Omega |\Phi\rangle = \epsilon_{\nu\mu} \psi_{-1/2}^\mu \tilde{\psi}_{-1/2}^\nu |0\rangle_{-1} \otimes |\tilde{0}\rangle_{-1} \quad (2.21)$$

so that only the states with symmetric polarizations are invariant under Ω . Thus, only the graviton and the dilaton survive the Ω projection while the antisymmetric Kalb-Ramond field is removed.

Now, let us consider the massless states of the R-R sector which are represented by

$$|\Psi\rangle = \frac{1}{(p+2)!} F_{\mu_0 \dots \mu_{p+2}} (C\Gamma^{\mu_0 \dots \mu_{p+2}})_{\alpha\beta} |S^\alpha\rangle_{-1/2} \otimes |\tilde{S}^\beta\rangle_{-1/2} \quad (2.22)$$

where C is the charge conjugation matrix. This state is physical if the antisymmetric polarization tensor satisfies $dF = d^*F = 0$ which are precisely the Bianchi identities and the Maxwell equations for a field strength. Then, using Eqs. (2.17) and (2.19), we easily find

$$\Omega|\Psi\rangle = -\frac{1}{(p+2)!} F_{\mu_0 \dots \mu_{p+2}} (C\Gamma^{\mu_0 \dots \mu_{p+2}})_{\beta\alpha} |S^\alpha\rangle_{-1/2} \otimes |\tilde{S}^\beta\rangle_{-1/2} \quad (2.23)$$

so that only those states for which

$$(C\Gamma^{\mu_0 \dots \mu_{p+2}})^T = - (C\Gamma^{\mu_0 \dots \mu_{p+2}}) \quad (2.24)$$

are even under Ω and must be kept. A little algebra shows that Eq. (2.24) is true if $p = 1, 5, 9$, so that the only R-R potentials that survive the Ω projection are a 2-form, its magnetic dual 6-form and a non-dynamical 10-form. Consequently, in the Type I theory only D1, D5 and D9-branes can exist as (isolated) brane configurations that carry R-R charges.

As we have mentioned before, the closed string states which are invariant under Ω do not represent the whole content of the theory. In fact, in the orientifold construction one can also impose twisted boundary conditions on the string coordinates, that is one can require that they be periodic up to an Ω transformation. In our case this amounts to impose that

$$X^\mu(\tau, \sigma) = \Omega X^\mu(\tau, \sigma - \pi) \Omega^{-1} = X^\mu(\tau, -\sigma) \quad , \quad (2.25)$$

where in the last step we have used the explicit definition (2.14). It is easy to realize that the twisted boundary condition (2.25) enforces an identification between the left and the right-moving parts since it implies that $\alpha_n^\mu = \tilde{\alpha}_n^\mu$. Thus, the string coordinates which satisfy Eq. (2.25) and which we denote by X_N^μ , have the same mode expansion as the coordinates of the *open* string with Neumann boundary conditions, namely

$$X_N^\mu(\tau, \sigma) = x^\mu + (2\alpha')p^\mu \tau + i\sqrt{2\alpha'} \sum_{n \neq 0} \frac{\alpha_n^\mu}{n} e^{-in\tau} \cos n\sigma \quad . \quad (2.26)$$

Then, from Eq. (2.14) it is immediate to see that the action of the world-sheet parity on the oscillators of X_N^μ is

$$\Omega \alpha_n^\mu \Omega^{-1} = e^{in\pi} \alpha_n^\mu \quad (2.27)$$

Similarly, one finds that the fermionic oscillators of the open sector have the following transformation rule

$$\Omega \psi_r^\mu \Omega^{-1} = e^{ir\pi} \psi_r^\mu \quad (2.28)$$

To complete our description, we have to specify also the action of Ω on the ground states of the open sector. For simplicity and in analogy with what we have done before, we choose the (-1) superghost picture for the NS sector and the $(-1/2)$ superghost picture for the R sector [31]. Then, in an obvious notation, we posit that

$$\Omega |0\rangle_{-1} = -i |0\rangle_{-1} \quad (2.29)$$

and

$$\Omega |S^\alpha\rangle_{-1/2} = - |S^\alpha\rangle_{-1/2} \quad (2.30)$$

With this choice the massless states of both the NS and R sectors are odd under Ω and thus are removed from the spectrum. This is a standard result for unoriented open strings *without* Chan-Paton factors. On the contrary, if at the endpoints of the open strings we introduce Chan-Paton labels for the group $SO(N)$, then the NS and R massless states can survive the Ω projection. In this case, then, the spectrum will contain also $SO(N)$ vector gauge fields and their superpartners. However, this theory is consistent only for a particular choice of the gauge group. In fact, the cancellation of gauge anomalies in the low-energy effective action requires that $N = 32$ [33]. This condition can be understood also as a massless tadpole cancellation condition in the Type I string [34].

This construction can be also formulated in the modern language of D-branes and orientifold planes. In this context, one starts again with Type IIB theory and, in order to enforce the world-sheet parity projection, one puts in the background the so-called crosscap state (or orientifold 9-plane, $O9$) [34, 35]. This is a space-time filling object that is not dynamical, since no open strings can end on it, change its shape or switch on a gauge field. However, the crosscap state is a source of closed strings because it can emit, for example, dilatons, gravitons or a R-R 10-form potential ⁵. In order to cancel the tadpoles associated to these massless states,

⁵In units where the R-R charge of a D9-brane is +1, the R-R charge of the $O9$ -plane is -32 .

one is forced to modify the definition of the background of the theory by adding 32 D9-branes. This is clearly equivalent to introduce a sector of open strings with Neumann boundary conditions. Moreover, since there are 32 different D9 branes on which the open strings can terminate, it is as if their end-points carry 32 different charges, thus inducing the $SO(32)$ gauge group.

As we have mentioned before, also D1 and D5 branes are compatible with the orientifold projection (or equivalently with the presence of the orientifold $O9$ -plane). Therefore, besides the sectors previously considered, the Type I theory contains also other sectors corresponding to open strings defined on these D-branes. Since in particular the open strings living on the D1 branes of Type I are relevant for our later discussion, we now discuss their main properties.

Let us consider a supersymmetric D1 brane of Type I, and the open strings that live on its two-dimensional world-sheet. These strings, called 1-1 strings since they have both end-points on the D1 brane, are characterized by two longitudinal coordinates $X_N^a(\tau, \sigma)$ with Neumann boundary conditions ($a = 0, 1$) and by eight transverse coordinates $X_D^i(\tau, \sigma)$ with Dirichlet boundary conditions ($i = 2, \dots, 9$). The longitudinal coordinates have the usual mode expansion as in Eq. (2.26), while for the transverse coordinates one has

$$X_D^i(\tau, \sigma) = y^i + \sqrt{2\alpha'} \sum_{n \neq 0} \frac{\alpha_n^\mu}{n} e^{-in\tau} \sin n\sigma \quad (2.31)$$

where y^i denotes the position of the D1 brane in the transverse space⁶. Consequently, the action of Ω on the “Dirichlet” oscillators is different from the one on the “Neumann” oscillators which we have discussed in the previous subsection. Indeed, using Eqs. (2.14) and (2.31), it is easy to find that

$$\Omega \alpha_n^i \Omega^{-1} = -e^{in\pi} \alpha_n^i \quad (2.32)$$

to be compared with Eq. (2.27). World-sheet supersymmetry requires that the action of Ω on the fermionic oscillators in the Dirichlet directions be given by

$$\Omega \psi_r^i \Omega^{-1} = -e^{ir\pi} \psi_r^i \quad (2.33)$$

for both the NS and R sectors. Finally, on the ground states of the 1-1 strings one has [3]

$$\Omega |0\rangle_{-1} = -i |0\rangle_{-1} \quad (2.34)$$

⁶More generally, if an open string is stretched between two D-branes located respectively at y_1^i and y_2^i , the first term in Eq. (2.31) is replaced by $y_1^i + \left(\frac{y_2^i - y_1^i}{\pi}\right) \sigma$.

for the NS sector in the -1 superghost picture, and

$$\Omega|S^\alpha\rangle_{-1/2} = -\Gamma^2 \cdots \Gamma^9 |S^\alpha\rangle_{-1/2} \quad (2.35)$$

for the R sector in the $-1/2$ superghost picture. Eq. (2.34) is identical to the corresponding one for the purely Neumann strings since the NS vacuum, being a Lorentz scalar, is insensitive to the extra minus sign that Ω picks up in the Dirichlet directions. On the contrary, the R vacuum is a (chiral) spinor, and the extra minus sign in the action of Ω on the transverse fermions $\psi^2, \dots, \psi^9$ is not irrelevant, and indeed it translates into the particular structure of Eq. (2.35).

Using these rules, one can easily find that the spectrum of the 1-1 open strings of Type I contains eight massless states both in the NS and in the R sector. The massless modes of the NS sector correspond to a vector that accounts for the freedom of translating the D1 brane along its eight transverse directions. The massless states of the R sector, instead, correspond to a chiral spinor χ such that $\Gamma^2 \cdots \Gamma^9 \chi = -\chi$, as it follows from Eq. (2.35). Upon quantization, the eight components of χ account for the $2^{8/2} = 16$ degeneracy of the D1 brane of Type I, as it should be expected for a BPS configuration of the $N = 1$ supersymmetry algebra in ten dimensions.

We conclude by mentioning that in the Type I theory there exist also open strings with mixed boundary conditions: Neumann at one end-point and Dirichlet at the other one. Strings of this kind are for example the 1-9 open strings that are suspended between a D1 brane and one of the D9-branes of the Type I background, or the 9-1 strings which have the opposite orientation. These strings are characterized by eight directions with mixed boundary conditions, and the states in their spectrum can not be eigenvectors of Ω since this operator transforms a Neumann-Dirichlet string into a Dirichlet-Neumann one. In this case, then, the orientifold projection simply requires to keep all symmetric combinations of 1-9 and 9-1 configurations.

3 The Type I theory and its D1 brane

The $SO(32)$ heterotic string and the Type I theory discussed in the previous section look so different from each other that they seem completely unrelated. The only evident similarity between them is that they have the same massless field content.

However, this fact could be interpreted simply as an accident due to supersymmetry, because in ten dimensions the invariance under local supersymmetry completely determines the form of the action for gauge fields coupled to gravity. This interpretation seems to be further confirmed by the fact that any resemblance between the spectra of the two theories disappears as soon as one goes to the first excited level. For instance, we have mentioned that in the heterotic theory the states of the first mass level do not always transform in the same way under the gauge group, but can be scalars, spinors, second-rank symmetric traceless tensors, second-rank or fourth-rank antisymmetric tensors of $SO(32)$ [1]. On the contrary, in the Type I theory the color degrees of freedom are encoded through the Chan-Paton factors (or through the 32 D9-branes of the Type I background in the modern language) and thus all perturbative states of this model transform in the adjoint representation of $SO(32)$. This is a standard result that occurs in any string theory where the gauge group is introduced by means of the Chan-Paton procedure.

However, despite the appearance, we nowadays know that the $SO(32)$ heterotic string and the Type I theory describe the same physics [3, 36], and are actually the two members of a dual pair of theories related to each other in a non-perturbative way. The first hint of this equivalence comes from the structure of their low-energy effective actions. Focusing for example on the kinetic term of the gauge fields, one has

$$S_H \sim \int d^{10}x \left[\sqrt{-\det(G_H)} e^{-2\phi_H} \text{Tr}(F_H^2) + \dots \right] \quad (3.1)$$

for the heterotic string, and

$$S_I \sim \int d^{10}x \left[\sqrt{-\det(G_I)} e^{-\phi_I} \text{Tr}(F_I^2) + \dots \right] \quad (3.2)$$

for the Type I theory. In these expressions, G denotes the (string frame) metric, ϕ the dilaton field, and F the $SO(32)$ field strength. The two actions (3.1) and (3.2) can be mapped into each other by a field redefinition that involves a change of the metric

$$G_{\mu\nu}^H \leftrightarrow e^{-\phi_I} G_{\mu\nu}^I \quad (3.3)$$

and a change of the dilaton

$$\phi_H \leftrightarrow -\phi_I \quad (3.4)$$

Since e^ϕ is the string coupling constant, Eq. (3.4) implies that if there has to be any relation between the heterotic and the Type I strings, then the strong coupling of one theory has to be connected to the weak coupling of the other. This fact also explains why the two perturbative spectra are so different; indeed, by increasing

the value of the coupling constant, a generic perturbative heterotic state becomes highly unstable and is not expected to appear in the Type I theory, and viceversa.

However there are some string configurations that can not decay; for this reason, they are expected to be present in both theories at all values of the coupling constant, in particular in the strong coupling region of one theory or, equivalently, in the weak coupling regime of the dual theory. Finding such stable states and checking their multiplicities is therefore a test of the duality conjecture that goes well beyond the arguments based on the structure of the effective actions. A well known class of such stable configurations consists of the BPS states. The main feature of these states is that they are annihilated by some supersymmetry generators so that they carry a charge under one of the gauge fields present in the theory and possess a mass which is completely determined by this charge. Furthermore, these BPS configurations are protected from quantum radiative corrections, and thus it is particularly simple to follow their fate when the coupling constant is increased. For instance, a heterotic string wrapped around a compact dimension breaks half the original supersymmetries and is charged under the antisymmetric two-form present in the gravitational multiplet, the charge being simply its winding number. This is a BPS configuration and hence it should appear in the Type I theory as a supersymmetric soliton. The natural candidate for this is the D1 brane of Type I, since one expects that the gravitational two-form gauge field is mapped in the R-R one, just like in the strong/weak duality of Type IIB string. Since we are considering BPS configurations, we can immediately test this proposal. In fact, if it is correct, the tension

$$\tau_H = \frac{1}{2\pi\alpha'} \quad (3.5)$$

of the fundamental heterotic string should exactly match the tension

$$\tau_{D1} = \frac{1}{2\pi\alpha'g} \quad (3.6)$$

of the D1 brane of Type I ⁷. This is indeed what happens if we use Eq. (3.3) to set the same scale in both theories.

The identification between the D1 brane of Type I and the fundamental string of the heterotic theory can be tested at a deeper level; in fact it is possible to show that the world-sheet structure of a (wrapped) heterotic string is exactly reproduced by the world-sheet dynamics on a (wrapped) D1 brane [3]. Here we recall this argument. In the presence of a D1 brane, the space-time is naturally divided into

⁷These tensions are written using the normalizations and units of Type II theories.

a two-dimensional Minkowski part along the world-sheet of the D1 brane (spanned by x^0 and x^1), and an eight-dimensional Euclidean part transverse to it (spanned by $x^2, \dots, x^9$). Consequently, the Lorentz group $SO(1, 9)$ is broken into $SO(1, 1) \times SO(8)$. The space transverse to the D1 brane is going to be identified with the space where the heterotic string is embedded in the light-cone gauge, while the world-sheet of the D1 brane is going to play the role of the world-sheet of the heterotic string. To see this, we will first consider the lightest states of the 1-1 open strings attached to the D1 brane which are responsible for its dynamics, and then identify the corresponding massless fields with the various string coordinates of the heterotic theory. In the NS sector we can construct the following massless states

$$\psi_{-1/2}^{\mu} |k\rangle_{-1} \quad (3.7)$$

with $k^2 = 0$. Since the momentum k is only along the longitudinal directions, the field associated to this state will depend only on x^0 and x^1 . When the index μ is either 0 or 1 (*i.e.* in a NN direction), the state (3.7) is odd under Ω (see Eqs. (2.28) and (2.29)), and thus it is not present in the Type I theory. This means that on the D1 brane of Type I there are no massless vectors, and thus no local gauge symmetry. On the contrary, when the index μ is $2, \dots, 9$ (*i.e.* in a DD direction) the state (3.7) is even under Ω (see Eqs. (2.33) and (2.34)), and thus survives the orientifold projection. The corresponding fields, $\phi^i(x^0, x^1)$ with $i = 2, \dots, 9$, are therefore scalars with respect to the world-sheet group $SO(1, 1)$ but are vectors with respect to the transverse group $SO(8)$. These are the same features of the embedding coordinates $X^i(\tau, \sigma)$ of the heterotic string, which therefore can be identified with $\phi^i(x^0, x^1)$ provided that we interpret x^0 and x^1 as τ and σ respectively. Notice since σ varies between 0 and π , we deduce that the D1 brane has to be wrapped on a radius R so that $x^1/2R$ can play the role of σ .

Let us now consider the massless states in the R sector of the 1-1 strings which are given by the spinor vacua

$$|S^{\alpha}; k\rangle_{-1/2} \quad (3.8)$$

with $k^2 = 0$. The field associated to these states is a chiral spinor χ obeying the massless Dirac equation

$$\not{k}\chi = (k_0\Gamma^0 + k_1\Gamma^1)\chi = 0 \quad (3.9)$$

However not all chiral spinors are acceptable in the Type I theory. In fact, as we have seen in Section 2.2, the orientifold projection acts on the fermionic zero-modes of the 1-1 strings as $\Omega = -\Gamma^2 \dots \Gamma^9$, and thus, in order to be even under Ω , the

spinor χ has to satisfy the further constraint

$$\Gamma^2 \dots \Gamma^9 \chi = -\chi \quad (3.10)$$

This condition tells us that in the decomposition of the chiral spinor representation of $SO(1, 9)$ into $(+\frac{1}{2}, \mathbf{8}_s) \oplus (-\frac{1}{2}, \mathbf{8}_c)$, only the second term is selected by the Ω projection. Notice that since χ is chiral, Eq. (3.10) implies that

$$\Gamma^0 \Gamma^1 \chi = -\chi \quad (3.11)$$

Hence, from the Dirac equation (3.9) we deduce that $k_0 = -k_1$, *i.e.* χ is a right-moving spinor in the $\mathbf{8}_c$ of $SO(8)$. These are precisely the same features of the fermionic coordinates of the heterotic string in the Green-Schwarz formulation!

In order to complete our description, we still have to reproduce in the D1 brane context the color degrees of freedom that in the heterotic theory are represented by the 32 left-moving fermions resulting from the fermionization of the 16 internal coordinates. On the Type I side these degrees of freedom are provided by the 1-9 and 9-1 open strings that stretch between the D1 brane and the 32 D9-branes of the background. The NS sector of these open strings does not contain any massless excitation, since any physical state in this sector has to satisfy the Virasoro constraint

$$\left(L_0 + \frac{1}{2}\right)|v\rangle = 0 \quad . \quad (3.12)$$

Notice that in this equation we have used the value $a = +1/2$ of the NS intercept for open strings with eight directions with mixed boundary conditions⁸. Hence in this NS sector even the ground state is massive. On the contrary, in the R sector the world-sheet supersymmetry is preserved by the boundary conditions and the intercept is always vanishing. Therefore, the R ground states are always massless. To see which are the other features of these states in the 1-9 and 9-1 sectors, let us recall that the fermions ψ^0 and ψ^1 (which carry an index in a NN direction) have the standard integral mode expansion and possess zero-modes. Instead, the fermions $\psi^2, \dots, \psi^9$ (which carry an index in a direction with mixed boundary conditions) have a half-integral mode expansion and do not possess zero-modes. Thus, the R ground states of these open strings are associated to a field Λ which is a spinor of

⁸Remember that the value of the intercept a can be easily obtained by summing the zero-point energy of all the light-cone coordinates. For the usual (*i.e.* all NN directions) NS sector one obtains $a = -1/2$, but when there are ν directions with mixed boundary conditions one gets $a = -1/2 + \nu/8$.

$SO(1,1)$ and a scalar of $SO(8)$. Furthermore, the GSO projection requires that such a spinor be chiral, *i.e.*

$$\Gamma^0 \Gamma^1 \Lambda = \Lambda \quad (3.13)$$

Combining this condition with the massless Dirac equation

$$\not{k} \Lambda = (k_0 \Gamma^0 + k_1 \Gamma^1) \Lambda = 0 \quad (3.14)$$

we deduce that $k_0 = k_1$, *i.e.* Λ is a left-moving world-sheet spinor! Notice that since x^1 is compact, the momentum k_1 in Eq. (3.8) must be an integer n in units of $1/R$.

As a matter of fact, this is not enough because we have also to specify whether we are considering the 1-9 or the 9-1 sector, and distinguish among the 32 different D9-branes of the background. This information can be provided by attaching to the state in Eq. (3.8) appropriate Chan-Paton factors, which we denote by λ_{19}^I ($I = 1, \dots, 32$) for the 1-9 sector, and by λ_{91}^I ($I = 1, \dots, 32$) for the 9-1 sector. Since the twist operator Ω exchanges the string orientation, it converts a 1-9 string into a 9-1 string and viceversa, *i.e.*

$$\Omega : \lambda_{19}^I \leftrightarrow \lambda_{91}^I \quad (3.15)$$

Therefore, the ground state which is selected by the Ω projection is given by the following symmetric combination

$$\lambda_{19}^I |S^+; n\rangle_{-1/2} + \lambda_{91}^I |S^+; n\rangle_{-1/2} \quad (3.16)$$

By considering all possible values for the momentum (that is by summing over n), we can construct a left-moving spinor with an index I in the vector representation of $SO(32)$ and with the following mode expansion

$$\Lambda^I(\tau, \sigma) = \sum_n \Lambda_n^I e^{i2n(\tau+\sigma)} \quad (3.17)$$

This is precisely the expansion of the 32 heterotic fermions with *periodic* boundary conditions. Upon quantization, these periodic fermions give rise to the spinorial representations of $SO(32)$ (*i.e.* those representations with half-integer p^A in the notation of Section 2.1). In particular, using only the zero-modes Λ_0^I , we can construct the two representations $\mathbf{2}^{15}$ and $\mathbf{2}^{15'}$ of $SO(32)$ that correspond to $p^A = \pm 1/2$ for all A with an even or odd number of positive signs respectively. As we have seen in Section 2.1, the representation $\mathbf{2}^{15}$ appears at the first massive level of the

heterotic string. To obtain the remaining representations of $SO(32)$, it is necessary to construct 32 fermions with anti-periodic boundary conditions. To see how this is possible, we need to understand better the fate of the gauge fields on the D1 brane of Type I.

We have already seen that the orientation projection forbids to put a local gauge field on a single D1 brane; however, it is still possible that a *global* gauge field $A = \frac{\theta}{2\pi R}$ be compatible with Ω . This constant gauge field A can always be set to zero, at least locally, by means of the gauge transformation

$$A \rightarrow A' = A + i\Lambda^{-1}\partial_1\Lambda \quad (3.18)$$

where $\Lambda = e^{i\frac{\theta x^1}{2\pi R}}$. However, even if A is a pure gauge, its presence is not completely trivial since it affects the global properties of the charged objects. In fact, a field Φ which carries a charge q under A and makes a loop around the compact dimension picks up a non-trivial phase due to the Aharonov-Bohm effect which is given by

$$W = \exp \left\{ -iq \oint dx^1 A \right\} = e^{-iq\theta} . \quad (3.19)$$

Of course, by gauge invariance, this same result has to be obtained also in the gauge where $A = 0$. This is indeed the case because under the gauge transformation (3.18), the charged field Φ changes as follows

$$\Phi \rightarrow \Phi' = e^{i\frac{q\theta x^1}{2\pi R}} \Phi \quad (3.20)$$

Thus, in the gauge $A = 0$, the field of a charged object is no longer periodic and acquires the non-trivial phase (3.19) when it is transported around the compact dimension. This fact can be rephrased by saying that its momentum k^1 is shifted according to

$$k^1 \rightarrow k^1 + \frac{q\theta}{2\pi R} \quad (3.21)$$

In our specific configuration the field A is localized on the D1 brane, and thus any string with at most one end-point on the D1 brane is charged under A . This is precisely the case of the 1-9 and 9-1 strings that we are considering. By convention we can take the charge to be $q = 1$ for the 1-9 strings and to be $q = -1$ for the 9-1 strings. As we mentioned before, the states in the R sector of these strings must always appear in linear combinations of the type

$$\lambda_{19}^I(\dots) |S^+; n\rangle_{-1/2} \pm \lambda_{91}^I(\dots) |S^+; m\rangle_{-1/2} , \quad (3.22)$$

where we have explicitly written only the dependence on the momentum along the compact dimension and understood all other information in the dots (the sign between the two terms in Eq. (3.22) depends on the Ω -parity of the oscillator content). In the absence of any gauge field only those combinations with $n = m$ are kept. However, when there is the constant gauge field A , things are slightly different. First of all, we have to shift the momenta of the charged states according to Eq. (3.21), and thus, instead of Eq. (3.22), we must consider the superposition

$$\lambda_{19}^I(\dots) |S^+; n + \theta/2\pi\rangle_{-1/2} \pm \lambda_{91}^I(\dots) |S^+; m - \theta/2\pi\rangle_{-1/2} \quad (3.23)$$

Since n and m are integers, it is clear that this combination can be an eigenstate of Ω only if $\theta = 0$ or π . Consequently, on the D1 brane of Type I there exists a Z_2 symmetry which is all what remains of the original $U(1)$ gauge symmetry of the D1 brane of Type IIB after performing the orientifold projection. When $\theta = \pi$, the massless fields associated to Eq. (3.23) are 32 world-sheet spinors Λ^I with the following mode expansion

$$\Lambda^I(\tau, \sigma) = \sum_n \Lambda_n^I e^{i(2n+1)(\tau+\sigma)} \quad (3.24)$$

This is precisely the expansion of the 32 heterotic fermions with *antiperiodic* boundary conditions. Upon quantization, these fermions give rise to the integral representations of $SO(32)$ (*i.e.* those representations with integer p^A in the notation of Section 2.1). Representations of this kind are for instance the singlet **1**, the vector **32** and the adjoint **496**.

To have the complete equivalence between the low-energy world-sheet dynamics of the wrapped D1 brane of Type I and the heterotic string, we still have to impose the appropriate GSO projection on the 32 heterotic fermions Λ^I of Eqs. (3.17) and (3.24) in order to remove, for example, the representations **2^{15'}** and **32** which can be constructed using the Λ^I 's but which do not appear in the heterotic theory at the massless level. This reduction can be achieved by using the idea of the orientifold projection presented in Section 2.2. In fact, on the D1 brane of Type I we can introduce a discrete transformation Ξ that acts only on the Chan-Paton factors of the open strings and is defined as follows: Ξ multiplies by -1 each time there is an open string ending on the D1 brane and acts trivially otherwise. Then, it is clear that gauge invariant states (*i.e.* states that are invariant under the Z_2 gauge group which survives the Ω projection) are left unchanged by Ξ . In view of these considerations, it appears natural to regard as physical excitations of the D-strings

only the ones which are obtained by gauge invariant superpositions of 1-9 and 9-1 strings; in other words, it appears natural to gauge the discrete symmetry Ξ . We have already encountered an example of a gauged discrete symmetry in Section 2.2 when we discussed the Ω projection and derived unoriented strings from the Type IIB theory. Here we can apply a similar procedure to gauge the discrete symmetry Ξ . The first step is to consider only those states of the 1-9 and 9-1 spectrum that are invariant under Ξ (the states in the 1-1 and 9-9 sectors are obviously all invariant under Ξ). For example, if we define the spinor representation $\mathbf{2}^{15}$ to be even under Ξ , then it is clear that the states in the other spinor representation $\mathbf{2}^{15'}$ are odd, since they contain one more fermionic zero-mode that brings a Chan-Paton factor λ_{19} or λ_{91} , and hence an extra $-$ sign under Ξ . Thus, the states of $\mathbf{2}^{15'}$ are gauged away and the physical degrees of freedom correspond only to the representation $\mathbf{2}^{15}$, just as the heterotic GSO projection requires. However, as we know very well, this is not the whole story, since in gauging a discrete symmetry one has also to consider the twisted sector. In fact, one can impose on the fields living on the D-string generalized periodicity conditions by exploiting the transformation Ξ , just as we did in Eq. (2.25) for Ω . In the present case, the only fields that can feel Ξ are those in the 1-9 and 9-1 sectors, *i.e.* the Λ^I 's. Thus, the twisted sector is characterized by

$$\Lambda^I(\tau, \sigma + \pi) = \Xi \Lambda^I(\tau, \sigma) \Xi^{-1} = -\Lambda^I(\tau, \sigma) . \quad (3.25)$$

These are precisely the antiperiodic boundary conditions satisfied by the expansion (3.24) which was obtained by introducing the Z_2 -Wilson line on the D-string. Therefore, one can say that introducing this Wilson line is equivalent to introducing the twisted sector of Λ^I under Ξ . Of course, also among the states of the twisted sector we have to select only the Ξ -invariant ones, which have an even number of Λ^I oscillators, since each one of them contributes with a $-$ sign to the Ξ parity. Then, one can check that, for example, the vector representation $\mathbf{32}$ is gauged away while the representations $\mathbf{1}$ and $\mathbf{496}$ are kept, thus reproducing the condition $\sum n_i = 2Z$ of the heterotic side (see Eq. (2.5)).

4 A Non-BPS D-particle in the Type I theory

Up to now, we have focused on BPS objects and exploited their stability to make a test of the non-perturbative duality between the $SO(32)$ heterotic string and the Type I theory. However, string theories often contain in their spectrum states that are *stable* but non-BPS, and thus one may wonder whether such states can provide the basis for duality tests that are less dependent on supersymmetry. Let us consider for instance a dual pair of theories A and B , and take all perturbative states of A that carry the same set of conserved quantum numbers $\{\alpha\}$. Clearly, the lightest ones among these states are absolutely stable, since any possible decay is energetically unfavorable, and exist for all values of the coupling constant of A . Hence, if the duality between A and B is correct, there should exist stable non-perturbative configurations of B with the quantum numbers $\{\alpha\}$. Finding such configurations and checking their multiplicities is therefore a significant test of the duality between A and B which does not rely on BPS or supersymmetry arguments. In this section we are going to apply this strategy and confirm the $SO(32)$ heterotic/Type I duality by using stable non-BPS objects. Then, in the following section we will put on more solid ground our results by providing an exact conformal field theory description of these non-BPS configurations.

We have encountered absolutely stable states when we studied the first excited level of the perturbative spectrum of the heterotic string: in fact, the massive states in the representation $\mathbf{2}^{15}$ of $SO(32)$ are the lightest ones transforming as spinors of the gauge group, since the massless level contains only states in the singlet and adjoint representations of $SO(32)$ (see Eq. (2.8)). Thus, if the heterotic/Type I duality is correct, we should find in the weakly coupled Type I theory some non-perturbative objects that behave as massive particles in the spinor representation of $SO(32)$. Finding such configurations and providing their explicit description will be our goals. As mentioned above, in the heterotic theory the lightest spinors of $SO(32)$ appear at the first massive level and their mass is $M = 2/\sqrt{\alpha'}$ at $g_H = 0$. As one increases the heterotic coupling constant, the mass M gets renormalized, since there are no constraints on it coming from supersymmetry, and so, for a general value of g_H , we can write

$$M = \frac{2}{\sqrt{\alpha'}} f(g_H) \tag{4.1}$$

where the function $f(g_H)$ can in principle be computed in perturbation theory and is such that $f(0) = 1$. By going to the Type I theory and using the duality relations

(3.3) and (3.4), we can rewrite the mass M as follows

$$M = \frac{2}{\sqrt{\alpha' g_I}} \tilde{f}(g_I) \quad (4.2)$$

where $\tilde{f}(g) \equiv f(1/g)$. Thus, contrarily to what happened for the tension of the D-string (3.6), the mass M of these stable non-BPS states for $g_I \rightarrow 0$ is not simply the translation of the heterotic mass at $g_H = 0$, but differs from it by a factor of $\tilde{f}(g_I \rightarrow 0)$. Of course, such a factor is not accessible to perturbation theory and must be determined using different methods.

Let us then examine more carefully the structure of these non-BPS configurations. On the heterotic side they are simply excited states of the fundamental string; thus, it is natural to expect that the D1 brane is involved on the Type I side. However a single D1 brane can not reproduce the features of our non-BPS states for several reasons. First of all, a D1 brane is a BPS configuration; moreover, as we have seen in the previous section, it is dual to a fundamental heterotic string with winding number 1. On the contrary, our non-BPS configurations correspond to *unwrapped* strings, *i.e.* to states that are not charged under the gravitational 2-form of the heterotic theory, and therefore they should be dual to Type I configurations that are neutral under the R-R 2-form potential. The simplest system constructed out of D1 branes but without any R-R charge, is clearly the superposition of a D1 brane and an anti-D1 brane ⁹. However, this configuration is *unstable* due to presence of a tachyonic mode arising from the open strings stretched between the brane and the anti-brane [14]. To see this, let us study the brane/anti-brane pair in more detail. When we put an anti-D1 brane on top of a D1 brane, we have four types of open strings:

- the 1-1 strings with both end-points on the D1 brane;
- the $\bar{1}\text{-}\bar{1}$ strings with both end-points on the anti-D1 brane;
- the 1- $\bar{1}$ strings starting from the D1 brane and ending on the anti-D1 brane;
- the $\bar{1}$ -1 strings starting from the anti-D1 brane and ending on the D1 brane.

As we have seen before, convenient way of encoding the information about the end-points of the open strings is by means of Chan-Paton factors. In the present case

⁹An anti-D p brane is simply a D p brane carrying a negative charge under the R-R $p + 1$ -form potential.

we can introduce 2×2 Chan-Paton matrices λ as follows

$$\text{for a } 1\text{-}1 \text{ string} \rightarrow \lambda_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad (4.3)$$

$$\text{for a } \bar{1}\text{-}\bar{1} \text{ string} \rightarrow \lambda_{\bar{1}\bar{1}} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad (4.4)$$

$$\text{for a } 1\text{-}\bar{1} \text{ string} \rightarrow \lambda_{1\bar{1}} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad (4.5)$$

$$\text{for a } \bar{1}\text{-}1 \text{ string} \rightarrow \lambda_{\bar{1}1} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad (4.6)$$

The $1\text{-}\bar{1}$ (and $\bar{1}\text{-}1$) strings differ in a crucial way from the $1\text{-}1$ strings considered in Section 2.2 because they are characterized by a “wrong” GSO projection, which, for example, selects the tachyonic ground state and removes the massless vectors in the NS sector. This fact can be understood more easily by studying this same system from the closed string point of view, where the interactions are seen as a tree-level exchange of closed string states along a cylinder between the brane and the anti-brane. However, for the sake of clarity, we first analyze what happens in the more familiar case of a pair formed by two D-branes on top of each other. The interaction between them is computed by summing the contributions to the exchange energy of all closed string states and can be organized as a sum of the even closed string spin structures: NS-NS, NS-NS $(-1)^F$ and R-R¹⁰. Of course, this is completely equivalent to compute the one-loop vacuum energy of open strings stretched between the two D-branes, which is organized as a sum of the even open string spin structures: NS, NS $(-1)^F$ and R. In fact, by means of a modular transformation $\tau_c \rightarrow \tau_a = 1/\tau_c$, it is possible to map the length τ_c of the cylinder described by the closed strings, into the modular parameter of the annulus τ_a spanned by the open strings. Then, using the modular properties of the Jacobi θ -functions that appear in the amplitude, one can explicitly check that under the above modular transformation the contribution coming from the exchange of R-R states is exactly mapped into the contribution of the NS $(-1)^F$ sector of the $1\text{-}1$ open strings; in other words, the exchange of R-R states between two D1 branes gives the same result that one finds by computing the one-loop vacuum energy of the NS sector of the $1\text{-}1$ open strings weighted with the GSO operator $(-1)^F$. The correspondence between the

¹⁰In the cases we are studying in this paper, the odd spin structure R-R $(-1)^F$ never gives a contribution and thus will not be considered.

closed string channel and the open string one can be established also for the other spin structures, and the results are summarized in the following table

closed string	open string
NS-NS	NS
NS-NS $(-1)^F$	R
R-R	NS $(-1)^F$

(4.7)

In particular, we easily see that the sum of the NS-NS and R-R sectors of the closed string is equivalent to the NS sector of the open string with the usual GSO projection. Thus, the NS sector of the 1-1 and $\bar{1}-\bar{1}$ strings contains only states that are *even* under $(-1)^F$; hence the tachyon ground state is not allowed ¹¹.

Let us now consider an anti-D brane on top of a D brane. In this case, from the closed string point of view, the NS-NS and the NS-NS $(-1)^F$ spin structures give exactly the same contributions as before, but the contribution of the R-R spin-structure changes sign, since it is proportional to the product of the R-R charges of the two branes. According to table (4.7), this fact implies that the contribution of the NS $(-1)^F$ spin structure of the open channel changes sign, so that the NS sector of the open strings stretched between a brane and an anti-brane has the “wrong” GSO projection. Thus, the NS sector of the 1- $\bar{1}$ and $\bar{1}$ -1 strings contains only states that are *odd* under $(-1)^F$; in particular, the two tachyonic states

$$|k\rangle_{-1} \otimes \lambda_{1\bar{1}} \quad \text{and} \quad |k\rangle_{-1} \otimes \lambda_{\bar{1}1} \quad (4.8)$$

with $k^2 = 1/2\alpha'$ are permitted. In conclusion, we can say that the requirement that a cylinder diagram can be reinterpreted as an annulus diagram through a modular transformation fixes the type of GSO projection that one has to perform on the various kinds of open strings stretched between two branes ¹². Since these are distinguished by the Chan-Paton factors (see Eqs. (4.3)-(4.6)), it is natural to introduce a generalized GSO projection to also take them into account. Let us then define the operator $(-1)^{\mathcal{F}}$ that acts like the usual $(-1)^F$ on the open string oscillators and ground states, and that acts on the Chan-Paton factors according to

$$(-1)^{\mathcal{F}} : \quad \lambda \rightarrow \sigma^3 \lambda \sigma^3, \quad (4.9)$$

¹¹As usual, the NS Fock vacuum is taken to be odd under $(-1)^F$.

¹²This is also what happens in the so called Type 0 strings which have recently attracted some attention in the literature [37].

where σ^i are the Pauli matrices. Obviously, λ_{11} and $\lambda_{\bar{1}\bar{1}}$ are even under $(-1)^{\mathcal{F}}$, while $\lambda_{1\bar{1}}$ and $\lambda_{\bar{1}1}$ are odd. Thus, our previous findings can be summarized by saying that the spectrum of the open strings suspended between two D branes, or between a D brane and an anti-D brane, contains only states that are *even* under the extended GSO operator $(-1)^{\mathcal{F}}$.

The considerations presented so far apply to any system formed by a brane and an anti-brane of Type II, and do not make any use of the twist operator Ω . Thus, one may wonder whether the presence of the tachyons (4.8) in the D1/anti-D1 pair can be avoided by going to the Type I theory. We can easily see, however, that this is not the case because the tachyon instability is not removed by the Ω projection. In fact, the situation is similar to the one described in Eq. (3.22) because also here the orientation projection selects only a linear combination of the two tachyonic states (4.8). In particular, since the oscillator content of the tachyons is even, the combination $(\lambda_{1\bar{1}} - \lambda_{\bar{1}1})$ is projected out and only the symmetric combination of the Chan-Paton matrices survives, so that the tachyon in the Type I theory is given by

$$|T\rangle = |k\rangle_{-1} \otimes (\lambda_{1\bar{1}} + \lambda_{\bar{1}1}) = |k\rangle_{-1} \otimes \sigma^1 . \quad (4.10)$$

At this point, one has to understand whether the instability due to the existence of a tachyonic state in the spectrum is an incurable problem of the system, or is just a consequence of the fact that we are expanding the tachyon field T around a false vacuum. Of course, only in this second case one may hope that the D1/anti-D1 pair could form a bound state that has something to do with the perturbative heterotic state we are looking for. To give an answer to this question one should compute the tachyonic potential $\mathcal{V}(T)$ and see whether or not it has a minimum. Unfortunately, our knowledge of this potential is very limited and thus only a qualitative analysis can be performed along these lines. Let us nevertheless examine what can be said in this regard. First of all, from Eq. (4.10) it is easy to realize that $\mathcal{V}(T)$ is an even function of T . In fact, let us suppose for a moment that $\mathcal{V}(T)$ contains a term like cT^{2n+1} ; then from it one can construct a non-vanishing amplitude with an odd number of external tachyons which has to be reproduced by some non-vanishing string diagram. However, all open string amplitudes are multiplied by the trace of the Chan-Paton factors of the external states, which in our case is zero: indeed $\text{Tr}[(\sigma^1)^{2n+1}] = 0$. Therefore, only even powers of T are allowed to appear in the tachyonic potential ¹³. Another thing that is known about $\mathcal{V}(T)$ is that the

¹³This is to be contrasted with the tachyonic potential of the bosonic string where also odd powers of T are allowed.

coefficient of the T^4 term is positive [14, 38] so that it is conceivable to think that there exist a minimum for $\mathcal{V}(T)$. We now assume that all higher order terms in the potential do not change the mexican-hat shape produced by the first two terms, and that $\mathcal{V}(T)$ has an absolute minimum for $T = \pm T_0$ (since the potential is an even function, the minima always come in pairs).

Given these assumptions, we can pose the following question: if we allow the tachyon to roll down at the values $\pm T_0$ and stabilize the system, is it possible that after tachyon condensation the D1/anti-D1 pair describes the Type I D-particle we are looking for? The answer to this question is no. In fact, as we have seen in the previous section, both the D1 and the anti-D1 branes of Type I carry the quantum numbers of the spinorial representation of the gauge group, so that the resulting bound state cannot be a spinor of $SO(32)$, which instead is the distinctive feature of the particle we are searching. On the other hand, it is quite natural to expect that a brane-antibrane pair annihilates into the vacuum just like a particle-antiparticle pair may do. Actually, this can happen only if the total energy of the system is vanishing; in our particular case this condition means that the negative energy density of the tachyonic condensate has to exactly cancel the positive energy density of the two D-strings, *i.e.*

$$\mathcal{V}(\pm T_0) + 2\tau_1 = 0 \quad , \quad (4.11)$$

where τ_1 is the D1 brane tension given in Eq. (3.6). We can rephrase the content of Eq. (4.11) by saying that when the tachyon condenses everywhere to $\pm T_0$ the entire system stabilizes in a “trivial” way and reaches a configuration that is indistinguishable from the supersymmetric vacuum into which the D1 and anti-D1 branes have annihilated¹⁴. While this scenario seems quite reasonable at first sight, at a deeper analysis it looks too radical and surprising since it implies that the only stable configuration in which the tachyon can condense is the vacuum. Luckily, as we shall see, there are other possibilities for tachyon condensation which lead to more interesting results.

The basic problems with the D1/anti-D1 pair we have considered so far are that both branes are spinors of the gauge group and that the resulting system after tachyon condensation can have the same quantum numbers of the vacuum. However, we know from the previous section that there is a simple way to avoid this situation. In fact, by compactifying on a circle of radius R the direction along

¹⁴In the literature there are other examples in which tachyon condensation restores space-time supersymmetry; see for example Ref. [39]

which the D-strings extend and by switching on a Z_2 Wilson line, we can remove the zero-modes of the 1-9 strings and make the D1 brane a scalar of $SO(32)$. Hence, a pair formed by a D1 brane and an anti-D1 brane with a Z_2 Wilson line (*e.g.* $\theta = 0$ on the D1 brane and $\theta = \pi$ on the anti-D1 brane) has the right quantum numbers to transform in the spinor representation of $SO(32)$ and is a good candidate to describe the Type I particle. At first sight, it seems unlikely that the presence of a Wilson line may change radically the fate of the D1/anti-D1 system; in fact, a constant gauge field usually does not modify the energy density of a configuration because its field strength is vanishing. However, in our case, the situation is quite different: since our system as a whole transforms like a spinor of $SO(32)$, the condensation of the tachyonic field can not restore the vacuum and a new stable configuration with non-zero energy can be realized. This is possible because the tachyonic state (4.10) has only one endpoint on the anti-D1 brane where $\theta = \pi$ so that the corresponding field T is charged under the Wilson line and feels the presence of a non vanishing θ . Therefore, the tachyonic condensate is different from the one obtained for the system without Wilson line and its contribution to the energy is not longer that of Eq. (4.11). What is happening is that the constant gauge field changes the energy density of the system through the modifications it induces on the charged field T which is present in the background. To see how this mechanism can work, it is easier to perform the gauge transformation (3.18) in order to “switch off” the Wilson line and deal with a non-interacting system. We have already shown that in this gauge the wavefunction of an object with charge q is no longer periodic, but according to Eq. (3.19) it acquires a phase $e^{-iq\theta}$ each time it is translated around the wrapped anti-D1 brane. Since in our case $\theta = \pi$ and $q = 1$, this simply means that the momentum k^1 of the state (4.10) is quantized in *half-integer* multiples of $1/R$ so that the tachyonic field T has the following mode expansion on the D-string world-sheet

$$T(x^0, x^1) = \sum_{n=-\infty}^{\infty} T_n(x^0) e^{i\frac{(n+\frac{1}{2})}{R}x^1} \quad (4.12)$$

From this expression we immediately see that

$$T(x^0, x^1 + 2\pi R) = -T(x^0, x^1) \quad (4.13)$$

i.e. T satisfies antiperiodic boundary conditions. The fact that a bosonic field is not periodic may seem strange; however this should not come as a surprise since this is the behavior of any charged field in the presence of a Wilson line, while only the neutral (*i.e.* gauge invariant) quantities remain periodic. Because of

Eq. (4.13), a system where T is everywhere constant is no longer acceptable and this shows clearly that the tachyonic condensate gets modified by the presence of the Z_2 Wilson line. In fact, what becomes allowed by the new boundary conditions of T is a configuration that interpolates between the two minima $\pm T_0$; thus, in this case, the tachyonic condensate corresponds to a kink solution T_{kink} that passes from $-T_0$ to T_0 as one goes around the compact dimension, namely

$$\begin{aligned} T_{\text{kink}}(x^1) &\rightarrow -T_0 & \text{for } x^1 \rightarrow -\pi R \\ T_{\text{kink}}(x^1) &\rightarrow +T_0 & \text{for } x^1 \rightarrow +\pi R \end{aligned} \quad (4.14)$$

Of course, the energy of this configuration depends on the exact shape of the tachyonic kink T_{kink} for all values of x^1 and, according to our previous arguments, it contributes to the mass M of the Type I particle as follows

$$M = \int dx^1 \left[\mathcal{V}(T_{\text{kink}}(x^1)) + 2\tau_{D1} \right] \quad (4.15)$$

Since the kink solution is not explicitly known, we cannot use this equation to evaluate M ; however we can easily extract from it the behavior of M for small values of the Type I coupling constant g_I . In fact, both the classical potential $\mathcal{V}(T)$ and the brane tension τ_{D1} scale with the usual factor of $1/g_I$ since they originate from a string calculation on a disk. Thus the mass M of the non-BPS particle is proportional to the inverse of the coupling constant, exactly as it happens for the supersymmetric D-particle of Type IIA theory. By using this information in Eq. (4.2), we can see that for $g_I \rightarrow 0$ the function $\tilde{f}(g_I)$ is given by

$$\tilde{f}(g_I) = \frac{c}{\sqrt{g_I}} \quad (4.16)$$

where c is a constant. In order to fix its value, we have to leave the field theory analysis and systematically study the conformal field theory of the open strings living on the D1/anti-D1 system. This is what we are going to do in the next section. However, before concluding this section, we would like to present one more consideration at a qualitative level: if one insists that c be a finite constant for all values of the radius R , then $T_{\text{kink}}(x^1)$ has to be equal to $\pm T_0$ almost everywhere and the core of the kink has to be concentrated in the x^1 direction around a point, say $\bar{x}_1$. These requirements ensure that the energy density in Eq. (4.15) is vanishing everywhere because of Eq. (4.11), except around the position of the tachyonic soliton. Thus one can interpret the energy of this system as the mass of a *localized* particle. Furthermore, the possibility to put the core of the kink in an arbitrary point $\bar{x}_1$

corresponds to freedom that the particle has to move along the x_1 direction. In the following sections we will show that this stable non-BPS particle of Type I is actually a D-particle.

5 Conformal field theory at the critical radius

So far we have not really exploited the presence of the compact direction around which the D-strings are wrapped and, in some sense, we have implicitly taken the $R \rightarrow \infty$ limit. In this section, on the other hand, we consider the D1/anti-D1 system at the particular radius

$$R_c = \sqrt{\frac{\alpha'}{2}} , \quad (5.1)$$

and study the conformal field theory of the open strings living on its world-volume. In particular, following Ref. [6] we show that the non-BPS configuration corresponding to a D1/anti-D1 bound state is a stable D-particle of Type I. Then, in the next section, we will consider the decompactification limit $R \rightarrow \infty$ and show that this D-particle is stable even for $R > R_c$. The critical radius R_c is special for several reasons: first of all, the effective (mass)² of the scalar state (4.10), which is tachyonic in the uncompactified space, is non-negative. In fact, from the constraint $(L_0 - 1/2)|T\rangle = 0$, we find that

$$M_T^2 = \frac{\left(n + \frac{1}{2}\right)^2}{R_c^2} - \frac{1}{2\alpha'} \geq 0 , \quad (5.2)$$

where we have taken into account that the momentum along x_1 is quantized in half-integer units of R_c because of the presence of the Z_2 Wilson line (see also Eq. (4.12)). Moreover, from Eq. (5.2) it is easy to see that the lightest excitations of T are those with $n = 0$ and $n = -1$ and that, for these two values, M_T^2 is exactly zero¹⁵. Thus, the conformal field theory at R_c possesses two marginal operators corresponding to these two massless scalar states, which, as we will see, play a crucial role in our discussion.

¹⁵From these observations we see that at the critical radius the field T does not represent a true tachyon any more; however, since it becomes tachyonic in the decompactification limit, T will be still called tachyon.

A second peculiar feature of the critical radius (5.1) is that at R_c the conformal field theory generated by the coordinates X^1 and ψ^1 admits several different representations. In fact, since a bosonic field X compactified on a circle of radius R_c and with Neumann boundary conditions is equivalent to a couple of real NS fermions, the (X^1, ψ^1) system is equally well described by three fermions ξ, η and ψ^1 where ξ and η are defined by ¹⁶

$$e^{\pm \frac{i}{\sqrt{2\alpha'}} X^1} \simeq \frac{1}{\sqrt{2}}(\xi \pm i\eta) \quad , \quad \eta \xi \simeq \frac{i}{2\alpha'} \partial X^1 \quad (5.3)$$

Since the three fermions ξ, η and ψ^1 are on equal footing, we can use again the bosonization formulas in order to regroup the three fields in a different way and form a new boson. For example, if we use ξ and ψ^1 , we can define a new bosonic field ϕ according to

$$e^{\pm \frac{i}{\sqrt{2\alpha'}} \phi} \simeq \frac{1}{\sqrt{2}}(\xi \pm i\psi^1) \quad , \quad \psi^1 \xi \simeq \frac{i}{2\alpha'} \partial \phi \quad , \quad (5.4)$$

whereas if we use η and ψ^1 , we can introduce yet another field ϕ' by means of

$$e^{\pm \frac{i}{\sqrt{2\alpha'}} \phi'} \simeq \frac{1}{\sqrt{2}}(\eta \pm i\psi^1) \quad , \quad \psi^1 \eta \simeq \frac{i}{2\alpha'} \partial \phi' \quad . \quad (5.5)$$

In Appendix A we give some further details of this equivalence and show that the new fields ϕ and ϕ' are bosons compactified on a circle of radius R_c ; however, in most cases, Eqs. (5.3)-(5.5) are sufficient to translate the physical quantities in the different pictures. Since all these descriptions are equivalent, one may wonder what is the advantage of using, for instance, ϕ and η as independent fields, instead of the more natural ones X^1 and ψ^1 . As we will show momentarily following Ref. [6], it turns out that if one describes the system using ϕ and η , it is possible to encode the tachyonic background explicitly in the conformal field theory. In order to see why this simplification occurs, let us focus on the two massless excitations of the “tachyonic” state (4.10), *i.e.*

$$|T_{\pm}\rangle = e^{\pm \frac{i}{\sqrt{2\alpha'}} X^1} |0\rangle_{-1} \otimes \sigma_1 \quad (5.6)$$

and look at the explicit form of their vertex operators $\mathcal{V}_{T_{\pm}}$ in the various representations. Using the bosonization formulas (5.3)-(5.5) it is easy to see that

$$\mathcal{V}_{T_{\pm}}^{(-1)} = e^{\pm \frac{i}{\sqrt{2\alpha'}} X^1} \simeq \frac{1}{\sqrt{2}}(\xi \pm i\eta) \simeq \left[\pm \frac{i}{\sqrt{2}} \eta + \frac{1}{2} \left(e^{\frac{i}{\sqrt{2\alpha'}} \phi} + e^{-\frac{i}{\sqrt{2\alpha'}} \phi} \right) \right] \quad , \quad (5.7)$$

¹⁶In this and in the following formulas we understand the normal ordering.

where for simplicity we have understood the superghost part and the Chan-Paton factor σ^1 . From these relations we immediately realize that the states $|T_{\pm}\rangle$ of Eq. (5.6) can also be written either as

$$|T_{\pm}\rangle = \frac{1}{\sqrt{2}}(\xi_{-\frac{1}{2}} \pm i\eta_{-\frac{1}{2}})|0\rangle_{-1} \otimes \sigma_1 \quad , \quad (5.8)$$

or as

$$|T_{\pm}\rangle = \left[\pm \frac{i}{\sqrt{2}} \eta_{-\frac{1}{2}}|0\rangle_{\phi} + \frac{1}{2} \left(\left|+\frac{1}{2}\right\rangle_{\phi} + \left|-\frac{1}{2}\right\rangle_{\phi} \right) \right] |0\rangle_{-1} \otimes \sigma_1 \quad , \quad (5.9)$$

where we have denoted by $|\ell\rangle_{\phi}$ the vacuum of ϕ with momentum ℓ . In particular, in the latter representation the combination

$$|\mathcal{T}\rangle \equiv \frac{1}{\sqrt{2}i}(|T_{+}\rangle - |T_{-}\rangle) = \eta_{-\frac{1}{2}}|0\rangle_{\phi}|0\rangle_{-1} \otimes \sigma_1 \quad (5.10)$$

exhibits a very simple form since it becomes formally identical to a massless vector state at zero momentum in the -1 picture with the ψ oscillator substituted by η . This is a first hint to the fact that the “tachyonic” field associated to $|\mathcal{T}\rangle$ can be probably treated in the same manner of a gauge field.

As we have seen in Section 3, a constant gauge field in the X direction can be easily incorporated in a conformal field theory: indeed, one can use a *free* fields, provided that the X -momentum of any charged state is shifted as indicated in Eq. (3.21). Therefore, our working hypothesis is that, at the critical radius R_c , the “tachyonic” background T_{kink} discussed in the previous section can be described by a Wilson line of the “gauge” field $\mathcal{T}$ associated to the state (5.10). In order to identify which is the bosonic coordinate related to η and write the Wilson line, it is sufficient to look at the vertex operator corresponding to $|\mathcal{T}\rangle$ in the 0-superghost picture. In the original (X^1, ψ^1) description, we know [31] that the picture changing operation transforms the “tachyon” vertex operators as follows

$$\mathcal{V}_{T_{\pm}}^{(-1)} = e^{\pm \frac{i}{\sqrt{2\alpha'}} X^1} \rightarrow \mathcal{V}_{T_{\pm}}^{(0)} = \pm i \psi^1 e^{\pm \frac{i}{\sqrt{2\alpha'}} X^1} ; \quad (5.11)$$

then, by using the bosonization formulas (5.3) and (5.4), we can easily obtain the (ϕ, η) description of $\mathcal{V}_{T_{\pm}}^{(0)}$, and finally get

$$\mathcal{V}_{\mathcal{T}}^{(0)} = \frac{i}{\sqrt{2\alpha'}} \partial\phi \otimes \sigma^1 \quad . \quad (5.12)$$

This is identical to the vertex of the usual gauge boson at zero momentum, where ϕ plays the role of the coordinate X . From this equation, it is clear that $\mathcal{V}_{\mathcal{T}}^{(0)}$

represents a marginal operator which can be used to modify the theory. Let us recall that at the critical radius R_c our original D1/anti-D1 pair is not unstable, since there are no tachyonic excitations on its world-volume; however, it is always possible to deform the conformal field theory by inserting any number of operators $\mathcal{V}_T^{(0)}$ without paying any energy. This amounts to introduce a Wilson line along ϕ which can be parametrized as follows

$$W(A) = e^{i \frac{A}{2\sqrt{2\alpha'}} \oint d\sigma \partial_\sigma \phi \otimes \sigma^1} \quad (5.13)$$

In a qualitative sense, the constant A is equivalent to the tachyon vacuum expectation value since it multiplies the “tachyon” vertex operator $\mathcal{V}_T^{(0)}$. As we explained in Section 3, the states that are charged under a Wilson line have a momentum which is shifted according to Eq. (3.21) in the gauge where the constant gauge field is zero. In our present case, since $\mathcal{V}_T^{(0)}$ is proportional to σ^1 , the charged states of the theory are those with Chan-Paton factors σ^2 and σ^3 . If we denote their charge by q , then the Wilson line (5.13) induces the following shift in their ϕ -momentum k_ϕ :

$$k_\phi \rightarrow k_\phi - \frac{Aq}{2\sqrt{2\alpha'}} = k_\phi - \frac{qA}{4} \frac{1}{R_c} \quad (5.14)$$

which implies that the Kaluza-Klein index along ϕ is $n - qA/4$ with n integer. For later convenience, it is useful to define the combinations $\tau^\pm = \sigma^3 \pm i\sigma^2$ that correspond to states with charge $q = \mp 2$ under σ^1 :

$$[\sigma^1, \tau^\pm] = \mp 2 \tau^\pm . \quad (5.15)$$

The states with Chan-Paton factors $\mathbb{1}$ and σ^1 are instead neutral ($q = 0$) under the Wilson line (5.13) and thus are not affected by the presence of a non-trivial tachyonic background.

For different values of A the conformal field theory on the world-volume of the D1/anti-D1 pair at the critical radius describes different physical systems which are always characterized by the same mass M . In fact, since at R_c there are no tachyonic modes, the D1 and the anti-D1 branes with a Z_2 Wilson line form a bound state at threshold whose mass is the integral over the circle of the energy density of the two constituents; thus

$$M = 2 (2\pi R_c) \tau_{D1} = \frac{\sqrt{2}}{\sqrt{\alpha'}} \frac{1}{g} = \frac{1}{\sqrt{\alpha'} g_I} . \quad (5.16)$$

The fact that M is proportional to $1/g_I$ tells us that this bound state is a D-particle, a result which was already anticipated in the previous section. Notice that M is

a factor of $\sqrt{2}$ bigger than the mass of the supersymmetric (and BPS) D0-brane of Type IIA theory. From Eq. (5.16) we can also fix the value of the constant c defined by Eq. (4.16) finding $c = 1$. This is clearly a result which is not achievable using perturbative methods! We remark, however, that it can be obtained using the boundary state formalism as recently shown in Ref. [40].

As long as one studies the system at the critical radius R_c , all values of A are equally acceptable and lead to a consistent theory; however in the following section we will show that $A = 1$ is a preferred value because only for $A = 1$ it is meaningful to take the decompactification limit $R \rightarrow \infty$. Moreover, it turns out that at $A = 1$ the mass M of the system does not depend on the radius¹⁷. This is precisely the result that one would intuitively expect after tachyon condensation since in that case the D1/anti-D1 system should really describe a particle. However, also in the analysis at the critical radius it is clear that the case $A = 1$ is special and that only for this value of A the system exhibits the properties required for describing the stable non-BPS particle we are looking for.

A first peculiar feature of our system at $A = 1$ emerges from a detailed analysis of the open strings that live on the world-volume of the D1/anti-D1 pair. In order to see this, let us first consider the vacuum energy of the open strings in the original system (*i.e.* $A = 0$) which is defined as

$$\Gamma_{A=0} = 2V \int_0^\infty \frac{dt}{2t} \left[\text{Tr}_{\text{NS}} \left(\frac{1 + (-1)^{\mathcal{F}}}{2} q^{2L_0 - 1} \right) - \text{Tr}_{\text{R}} \left(\frac{1 + (-1)^{\mathcal{F}}}{2} q^{2L_0} \right) \right] \quad (5.17)$$

where V is the (infinite) length of the time direction and $q = e^{-t}$. The traces in Eq. (5.17) are to be computed on the NS and R sectors by taking into account the (extended) GSO projection that acts both on the string oscillators and on the Chan-Paton factors according to Eq. (4.9). In our present case there is one more thing to take into account, namely the existence of the Z_2 Wilson line on the anti-D1 brane. This implies that the string states with the diagonal Chan-Paton factors $\mathbb{1}$ and σ^3 (which have the standard GSO projection on the fermionic oscillators) are neutral under the Z_2 gauge field and their momentum in the compact direction X^1 is quantized in integer units of $1/R_c$. On the contrary, the states with the off-diagonal Chan-Paton factors σ^1 and $i\sigma^2$ (which have the “wrong” GSO on the fermionic oscillators) carry a Z_2 charge and thus their momentum along X^1 is quantized in half-integer units of $1/R_c$. This information can be encoded automatically in the

¹⁷Notice that this is the same behavior of the mass that we discussed at the end of the previous section.

formalism by introducing an operator h which corresponds to a $2\pi R_c$ translation along X^1 . Its explicit action on the string coordinates is

$$h : X^1 \rightarrow X^1 + 2\pi R_c \quad , \quad X^{\mu \neq 1} \rightarrow X^{\mu \neq 1} \quad , \quad \psi^\mu \rightarrow \psi^\mu \quad (5.18)$$

while it acts trivially as the identity on the NS and R ground states. If we define the action of h on the Chan-Paton factors λ by means of

$$h : \lambda \rightarrow \sigma^3 \lambda \sigma^3 \quad (5.19)$$

then it is not difficult to realize that only the states which are even under h belong to the physical spectrum and contribute to the vacuum energy Γ . Thus, the presence of the Z_2 Wilson line on the anti-D1 brane can be taken into account by introducing inside the traces of Eq. (5.17) the projector

$$\Pi_h = \frac{1+h}{2} \quad . \quad (5.20)$$

With this in mind, a standard calculation leads to

$$\begin{aligned} \Gamma_{A=0} = & 2V \int_0^\infty \frac{dt}{2t} (8\pi\alpha't)^{-1/2} \left\{ \sum_{n=-\infty}^\infty q^{4n^2} \left[\left(\frac{f_3(q)}{f_1(q)} \right)^8 - \left(\frac{f_4(q)}{f_1(q)} \right)^8 \right] \right. \\ & + \sum_{n=-\infty}^\infty q^{4(n+\frac{1}{2})^2} \left[\left(\frac{f_3(q)}{f_1(q)} \right)^8 + \left(\frac{f_4(q)}{f_1(q)} \right)^8 \right] \\ & \left. - \left[\sum_{n=-\infty}^\infty q^{4n^2} + \sum_{n=-\infty}^\infty q^{4(n+\frac{1}{2})^2} \right] \left(\frac{f_2(q)}{f_1(q)} \right)^8 \right\} \quad , \end{aligned} \quad (5.21)$$

where

$$\begin{aligned} f_1(q) &= q^{\frac{1}{12}} \prod_{n=1}^\infty (1 - q^{2n}) \quad , \quad f_2(q) = \sqrt{2} q^{\frac{1}{12}} \prod_{n=1}^\infty (1 + q^{2n}) \\ f_3(q) &= q^{-\frac{1}{24}} \prod_{n=1}^\infty (1 + q^{2n-1}) \quad , \quad f_4(q) = q^{-\frac{1}{24}} \prod_{n=1}^\infty (1 - q^{2n-1}) \quad . \end{aligned} \quad (5.22)$$

The structure of Eq. (5.21) is quite clear: apart from the factor of $(8\pi\alpha't)^{-1/2}$ coming from the integration over the continuum momentum k_0 along the time direction, the first line is the NS sector contribution of the open strings with diagonal Chan-Paton factors. The second line represents instead the contribution coming from the open strings with off-diagonal Chan-Paton matrices. Finally, the last line is the contribution of the R sector.

To perform this same calculation if $A \neq 0$, it is necessary to use the fields ϕ and η instead of X^1 and ψ^1 , since only in this way we know how to give the tachyon a non-vanishing vacuum expectation value in a conformal field theory framework. To obtain Γ_A it is convenient to start from the (ϕ, η) description of the system at $A = 0$ and select the physical states which are even both under $(-1)^{\mathcal{F}}$ and under h . In the (ϕ, η) representation these discrete symmetries are realized explicitly as reported in Tables 1, 2 and 3 of Appendix A. Then one can switch on the “tachyonic” Wilson line and follow how the various states of the spectrum are modified. In particular, the contributions coming from the NS open strings with Chan-Paton factors $\mathbb{1}$ and σ^1 are unchanged, since these two sectors are neutral under the Wilson line (5.13) and do not feel the presence of a non-trivial $\mathcal{T}$ background. Also the contribution of the R sector does not change when $A \neq 0$ because in this sector the field ϕ obeys mixed Neumann-Dirichlet boundary conditions so that its momentum is forced always to be zero (see Appendix A for some details). Thus, only the NS contribution of the strings with Chan-Paton factors $\tau^\pm$ depends on A . Putting all these things together, one finds that

$$\begin{aligned}
\Gamma_A = & 2V \int_0^\infty \frac{dt}{2t} (8\pi\alpha't)^{-1/2} \left\{ \frac{1}{2} \sum_{n=-\infty}^\infty q^{4n^2} \left[\left(\frac{f_3(q)}{f_1(q)} \right)^8 - \left(\frac{f_4(q)}{f_1(q)} \right)^8 \right] \right. \\
& + \frac{1}{2} \sum_{n=-\infty}^\infty q^{4(n+\frac{1}{2})^2} \left[\left(\frac{f_3(q)}{f_1(q)} \right)^8 + \left(\frac{f_4(q)}{f_1(q)} \right)^8 \right] \\
& + \frac{1}{2} \sum_{n=-\infty}^\infty \left[q^{4(n+\frac{A}{2})^2} \frac{(f_3(q))^8 - (f_4(q))^8}{(f_1(q))^8} + \sum_{n=-\infty}^\infty q^{4(n+\frac{1}{2}+\frac{A}{2})^2} \frac{(f_3(q))^8 + (f_4(q))^8}{(f_1(q))^8} \right] \\
& \left. - \left[\sum_{n=-\infty}^\infty q^{4n^2} + \sum_{n=-\infty}^\infty q^{4(n+\frac{1}{2})^2} \right] \left(\frac{f_2(q)}{f_1(q)} \right)^8 \right\} . \tag{5.23}
\end{aligned}$$

The A dependence is only in the third line which represents the contribution of the states with Chan-Paton factors $\tau^\pm$ whose charge under $W(A)$ is ∓ 2 (see Eq. (5.15)). In particular, the first term with an exponent proportional to $(n + \frac{A}{2})^2$, comes from the states with an odd number of η -oscillators, while the second term with an exponent proportional to $(n + \frac{1}{2} + \frac{A}{2})^2$, is produced by the states with an even number of η -oscillators.

At $A = 1$ a remarkable simplification occurs: all terms proportional to $f_4(q)$ cancel. This means that, for this value of A there is no NS $(-1)^F$ spin structure in the open string channel and consequently, according to Eq. (4.7), there is no R-R contribution to the vacuum energy from the closed string point of view ¹⁸! Thus,

¹⁸The cancellation of the R-R sector can also be seen directly in the closed string description

only for $A = 1$ the D1/anti-D1 pair really represents a neutral object under the R-R field and can be identified with the perturbative heterotic states with zero winding number.

Another important property of the system at $A = 1$ is that it becomes so tightly bound that it is not possible to separate the two constituent D1-branes without paying energy. To see this, let us consider the massless NS states

$$\psi_{-\frac{1}{2}}^i |0\rangle_{-1} \quad \text{with } i = 2, \dots, 9, \quad (5.24)$$

which describe the shape of a D-string, and see the modification induced on them by the tachyon background. Notice that actually we can construct two different states out of Eq. (5.24) by tensoring it with the Chan-Paton factors $\mathbb{1}$ or σ^3 (the other two Chan-Paton matrices σ^1 and $i\sigma^2$ are removed by the GSO projection). The first combination is neutral under the Wilson line $W(A)$ and thus remains massless for all values of A ; it corresponds to the marginal operators

$$\mathcal{V}^{i,(-1)} = \psi^i \otimes \mathbb{1} \quad (5.25)$$

for $i = 2, \dots, 9$ which account for the freedom of rigidly moving the D1/anti-D1 pair in the flat transverse directions X^i . On the other hand, the operator $\psi_{-\frac{1}{2}}^i \otimes \sigma^3$ represents a translation of the two branes in opposite directions and thus is related to the possibility of separating the two components of the system. However, being proportional to σ^3 , this type of state feels the effect of the tachyon background: if one introduces the matrices $\tau^\pm$, it is easy to find the modification induced by A , namely

$$\begin{aligned} \psi_{-\frac{1}{2}}^i |0\rangle_{-1} |0\rangle_\phi \otimes \sigma^3 &= \frac{1}{2} \psi_{-\frac{1}{2}}^i |0\rangle_{-1} |0\rangle_\phi \otimes (\tau^+ + \tau^-) \\ &\rightarrow \frac{1}{2} \psi_{-\frac{1}{2}}^i |0\rangle_{-1} \left(\left| -\frac{A}{2} \right\rangle_\phi \otimes \tau^+ + \left| \frac{A}{2} \right\rangle_\phi \otimes \tau^- \right). \end{aligned} \quad (5.26)$$

When $A \neq 0$ this state acquires a mass proportional to A^2 thus showing that there are no zero-modes permitting the separation of the D-string and the anti-D-string. Note that the r.h.s. of Eq. (5.26) is *not* invariant under the discrete transformation h corresponding to a $2\pi R_c$ translation along the compact direction. However, this is not surprising because we know from Section 4 that the periodicity of charged states is modified by the presence of a Wilson line: in our case, from Eqs. (5.14) and (3.19), we can see that a state of charge q picks up a phase equal to $e^{-i\frac{\pi A q}{2}}$ when it

by using the boundary state approach [40].

is transported once around the circle. Since in our normalizations all charged states have $|q| = 2$, we see that also the value $A = 2$ is particular: indeed, at $A = 2$ all states of the spectrum (even the charged ones!) have the same periodicity of the original theory at $A = 0$. Moreover, at $A = 2$ the vacuum energy (5.23) reduces to that of Eq. (5.21) confirming the fact that the conformal field theories at $A = 0$ and at $A = 2$ describe the same physical situation.

This analysis shows that, at the critical radius R_c , the presence of the “tachyonic” background can be nicely encoded in the conformal field theory by introducing a constant vacuum expectation value A for the field $\mathcal{T}$ (see Eqs. (5.10) and (5.12)). Moreover, A turns out to be an angular variable ranging from 0 to 2. The middle point of this interval ($A = 1$) has two special properties:

- at $A = 1$ the system as a whole does not emit any R-R field, so that it is possible to identify it with a heterotic state not charged under $B_{\mu\nu}$ at strong coupling;
- at $A = 1$ the D-string and the anti-D-string are most tightly bound, since the lightest state describing the separation of the two has maximum mass. In fact, when $A > 1$, one can consider a state similar to that of Eq. (5.26), but proportional to $\left(|1 - \frac{A}{2}\rangle_\phi \otimes \tau^+ + |-1 + \frac{A}{2}\rangle_\phi \otimes \tau^-\right)$; it has the right periodicity to be present in the physical spectrum and is lighter than the one of Eq. (5.26) with $A = 1$.

These observations suggest that the value $A = 1$ really represents the minimum of the tachyonic potential which corresponds to the kink solution discussed in the previous section. Thus, one expects that this value of A should give rise to a stable system also when we take the decompactification limit $R \rightarrow \infty$ and recover the full $SO(9)$ invariance. We note however, that even at the critical radius R_c it is possible to find a ninth zero mode corresponding to the freedom of moving the $\mathcal{T}$ soliton along the x^1 direction. Under a translation $x^1 \rightarrow x^1 + \epsilon$, the vertex operators (5.7) transform as

$$\delta_\epsilon \mathcal{V}_{T_\pm}^{(-1)} = \pm i \epsilon e^{\pm \frac{i}{\sqrt{2\alpha'}} X^1} \otimes \sigma^1 . \quad (5.27)$$

so that

$$\begin{aligned} \delta_\epsilon \mathcal{V}_T^{(-1)} &= \frac{1}{2i} \left(\delta_\epsilon \mathcal{V}_{T_+}^{(-1)} - \delta_\epsilon \mathcal{V}_{T_-}^{(-1)} \right) \\ &= \frac{\epsilon}{\sqrt{2}} \left(e^{+\frac{i}{\sqrt{2\alpha'}} X^1} + e^{-\frac{i}{\sqrt{2\alpha'}} X^1} \right) \otimes \sigma^1 \simeq \epsilon \xi \otimes \sigma^1 \end{aligned} \quad (5.28)$$

Thus, the marginal vertex operator

$$\mathcal{V}^{(-1)} = \xi \otimes \sigma^1 \quad (5.29)$$

represents the ninth zero mode we were looking for and can be combined with the eight vertices (5.25) to generate the $SO(9)$ symmetry associated to the freedom of moving the D-particle in space.

So far the twist operator Ω has not played any significant role. However, we conclude this section by stressing that the Ω projection is actually a crucial ingredient in the entire construction. In fact, even if it is possible to study the D1/anti-D1 pair in the context of Type IIB strings, one does not expect to find in this case a *stable* configuration corresponding to a D-particle. This is because Type IIB theory is self-dual under S-duality and there is no place for a stable non-BPS D-particle configuration. On the other hand, the instability of the D1/anti-D1 pair (with a constant gauge field $\theta = 1$) in the Type IIB theory is clearly displayed also at the level of the conformal field theory. In fact, if the worldsheet parity projection is not taken into account, one has to consider all open string states of the 1-1, $\bar{1}$ - $\bar{1}$, 1- $\bar{1}$ and $\bar{1}$ -1 sectors which are even under $(-1)^{\mathcal{F}}$ and h , without requiring that they be even under Ω . In our particular case, this implies that the “tachyonic” field $\mathcal{T}$ becomes complex, because besides the state in Eq. (4.10), there is also the combination with a Chan-Paton factor $i\sigma^2$. Moreover, the usual $U(1) \times U(1)$ gauge bosons of any brane/anti-brane pair with Chan-Paton factors $\mathbb{1}$ and σ^3 are also present in the spectrum of the oriented theory. At the critical radius R_c and at $A = 0$, these four massless states can be written in terms of η and ϕ according to

$$\begin{aligned} e^{\pm \frac{i}{\sqrt{2\alpha'}} X^1} |0\rangle_{-1} \otimes i\sigma^2 &\rightarrow \left[\pm \frac{i}{\sqrt{2}} \eta_{-\frac{1}{2}} |0\rangle_{\phi} + \frac{1}{2} \left(\left| \frac{1}{2} \right\rangle_{\phi} + \left| -\frac{1}{2} \right\rangle_{\phi} \right) \right] |0\rangle_{-1} \otimes i\sigma_2 , \\ \psi_{-\frac{1}{2}}^1 |0\rangle_{-1} \otimes \sigma^3 &\rightarrow \left[-\frac{i}{\sqrt{2}} \left(\left| \frac{1}{2} \right\rangle_{\phi} - \left| -\frac{1}{2} \right\rangle_{\phi} \right) \right] |0\rangle_{-1} \otimes \sigma_3 , \\ \psi_{-\frac{1}{2}}^1 |0\rangle_{-1} \otimes \mathbb{1} &\rightarrow \left[-\frac{i}{\sqrt{2}} \left(\left| \frac{1}{2} \right\rangle_{\phi} - \left| -\frac{1}{2} \right\rangle_{\phi} \right) \right] |0\rangle_{-1} \otimes \mathbb{1} . \end{aligned} \quad (5.30)$$

We remark that these states are odd under Ω and thus do exist in the Type I theory¹⁹. By taking a linear combination of the first two lines in Eq. (5.30) we

¹⁹It is worth pointing out that the action of the twist operator Ω on the NS vacuum of the 1- $\bar{1}$ and $\bar{1}$ -1 sectors is given by

$$\Omega |0\rangle_{-1} = |0\rangle_{-1}$$

i.e. there is an extra factor of i with respect to Eq. (2.29) which refers to the 1-1 sector. This fact can be understood using for example the arguments presented in Ref. [41]; see also Ref. [40].

obtain the following state

$$|\mathfrak{t}\rangle_{A=0} = \left(\left| \frac{1}{2} \right\rangle_{\phi} \otimes \tau^+ - \left| -\frac{1}{2} \right\rangle_{\phi} \otimes \tau^- \right) |0\rangle_{-1} . \quad (5.31)$$

The presence of the Chan-Paton factors $\tau^{\pm}$ shows that this state is charged under the tachyonic Wilson line $W(A)$. The other possible combinations that can be constructed out of Eq. (5.30) are instead neutral. In particular, if we set $A = 1$ which is the value corresponding to the tachyonic kink solution, the shift (5.14) of the ϕ -momentum transforms Eq. (5.31) into a very simple expression

$$|\mathfrak{t}\rangle_{A=1} = i |0\rangle_{\phi} |0\rangle_{-1} \otimes \sigma^2 . \quad (5.32)$$

It is not difficult to realize that this state has a *negative* mass², *i.e.* it is tachyonic! Thus, even if we have condensed the original tachyon field $\mathcal{T}$, the Type IIB system responds by creating a new tachyonic state. Thus, in Type IIB theory, there is an unavoidable tachyonic instability also at the critical radius whenever one gives a non-vanishing vacuum expectation value to the field $\mathcal{T}$. This should not be surprising because, as we have said before, the tachyon T of the Type IIB system is a complex field and the minimum of its potential does not consist any more of two separate points $(\pm T_0)$, but lies along a circle of radius $|T_0|$. Thus a kink configuration can not be stable because there is not any topological constraint that forbids its decay into the trivial solutions $T = T_0$ or $T = -T_0$. The possibility of this decay is precisely described by the tachyonic mode (5.32) that is present in the Type IIB theory, but not in the Type I model where it is projected out by Ω .

In conclusion we have shown that a D1/anti-D1 pair with a Z_2 Wilson line $\theta = \pi$ on the anti-brane and with a “tachyonic” Wilson line $A = 1$ at the critical radius R_c describes a D-particle which is non-BPS and unstable in Type IIB theory. However, by performing the Ω projection the tachyon instability is removed and the non-BPS D-particle of Type I is a stable configuration.

6 The Type I D-particle in Minkowski space

From the qualitative considerations of Section 4, it emerged that a the D1/anti-D1 pair with a Z_2 gauge Wilson line has the right quantum numbers to describe the $SO(32)$ spinor states which are present at the first massive level of the perturbative

heterotic spectrum. In the previous section we learned that at the critical radius $R_c = \sqrt{\alpha'/2}$ it is possible to describe exactly the tachyonic background living on the system and to study the conformal field theory of the open strings ending on it. Now it is crucial to show that, once the tachyon has condensed, the stability and the properties of the resulting bound state do not depend on the value of the radius and that it is possible to smoothly take the decompactification limit $R \rightarrow \infty$ to describe a D-particle of Type I in Minkowski space. To show this, we will proceed in the framework of perturbation theory: within the conformal field theory at critical radius $R = R_c$, we describe a small variation of R by inserting in the various amplitudes the operator $\mathcal{V}_R$ associated to the radius deformation. For instance, by calculating the one point function of the vertex operator (5.12), one can check the stability of our vacuum configuration; in fact, if the background satisfies the classical equations of motion all tadpoles are vanishing. The same amplitude with a $\mathcal{V}_R$ insertion describes the first order correction to the stability of this background arising from the $R_c \rightarrow R_c + \delta R$ variation. We have claimed that the value $A = 1$ is special, since it should represent a stable configuration also beyond the critical radius; if this is the case, it is clear that the tadpoles cancellation must not be effected by the variation of R and thus the amplitude

$$T_\phi = \langle T(\mathcal{V}_T(x) \mathcal{V}_R(z)) \rangle_A \quad (6.1)$$

should vanish for $A = 1$ ²⁰.

The explicit form of the marginal operator $\mathcal{V}_R$ is related to the graviton vertex at zero-momentum which, for instance, in the bosonic theory is given by

$$\mathcal{V}_R = \partial X(z) \bar{\partial} \tilde{X}(\bar{z}) . \quad (6.2)$$

The same result holds also for superstrings when $\mathcal{V}_R$ is written in the $(0,0)$ - superghost picture (that is, when it does not carry either left or right superghost charge), while, in the $(-1,-1)$ - picture, the bosonic coordinate X is substituted by ψ and the vertex reads

$$\mathcal{V}_R^{(-1,-1)} = \psi(z) \tilde{\psi}(\bar{z}) . \quad (6.3)$$

A second ingredient which is necessary for the computation of the amplitude (6.1) is the definition of the expectation value $\langle \dots \rangle_A$ in presence of the non trivial “tachyonic” background. From the analysis of the previous section we already know that in

²⁰Because of projective invariance, the correlation function in Eq. (6.1) does not depend on the positions of the vertex operators and it is convenient to exploit this fact by fixing $x = 1$ and $z = i$.

terms of the (η, ϕ) variables the “tachyonic” background can be easily taken into account by shifting the ϕ -momentum of charged states according to Eq. (5.14). However, this is *not* the only modification to disk amplitudes (like the one in Eq. (6.1)) induced by the presence of A because there is one more subtlety. In fact, one has to remember that in general, when closed string vertex operators are present, a gauge field Y couples also to the total winding number carried by the closed external states [42]. Thus, in the presence of a Y background, the expectation value on a disk reads as follows

$$\langle \dots \rangle_Y = \langle \dots e^{iY_I} \oint d\sigma^\alpha \partial_\alpha X^I \rangle . \quad (6.4)$$

Note that this modification is necessary to ensure the invariance of the disk amplitude under a T-duality that transforms the constant gauge field Y in the position of the D-brane. In our case, when the $\mathcal{T}$ background is written in the (η, ϕ) language, it has exactly the form of a constant gauge field and so, as Eq. (5.13) suggests, the expectation value $\langle \dots \rangle_A$ should be defined as in Eq. (6.4), namely

$$\langle \dots \rangle_A = \langle \dots \text{Tr} \left((\dots) e^{i \frac{A}{2\sqrt{2\alpha'}}} \oint d\sigma \partial_\sigma \phi \otimes \sigma^1 \right) \rangle , \quad (6.5)$$

where the Chan-Paton factors of the external open strings have to be inserted inside the trace. A final observation should be made about the time ordering T in Eq. (6.1): in order to avoid ambiguities among the open and closed vertex operators it is useful to perform the change of variables [43]

$$z \rightarrow z' = -\frac{z - i}{z + i} . \quad (6.6)$$

This transformation maps the upper half z -plane into the circle of unit radius and the lower part of the z -plane outside the same circle. Thus the closed string vertices are split and all the holomorphic parts are placed on the right, while all the antiholomorphic ones are placed on the left; the open string vertices are inserted in between, since they are forced to stay exactly on the unit circle.

Now we are in the position of evaluating the correlation function in Eq. (6.1); the first step is to translate the operator (6.3) in the (ϕ, η) language in order to be able to use the definition (6.5) for the expectation value in presence of a non trivial A . We have already performed this transformation at level of states (see Eq. (5.30)). Then, by repeating the same steps and exploiting the freedom to fix the punctures at 0, 1 and ∞ , it is easy to see that Eq. (6.1) can be explicitly rewritten as

$$T_\phi \sim \langle \left(e^{\frac{i}{\sqrt{2\alpha'}} \tilde{\phi}} - e^{-\frac{i}{\sqrt{2\alpha'}} \tilde{\phi}} \right) \Big|_\infty \partial\phi \Big|_1 \left(e^{\frac{i}{\sqrt{2\alpha'}} \phi} - e^{-\frac{i}{\sqrt{2\alpha'}} \phi} \right) \Big|_0 \text{Tr} \left(\sigma^1 e^{i \frac{A}{2\sqrt{2\alpha'}}} \oint d\sigma \partial_\sigma \phi \otimes \sigma^1 \right) \rangle , \quad (6.7)$$

where the left and the right fields ϕ and $\tilde{\phi}$ must be identified like in any disk amplitude. The structure of this equation is identical to that of a three-point function among open strings. So it is easy to see that two of the four terms arising from Eq. (6.7) are trivially zero because of ϕ -momentum conservation; in fact, as stressed in Appendix A, the new boson ϕ defined in Eq. (5.4) satisfies Neumann boundary conditions so that its momentum can not flow away from the disk boundary. The only two terms that give a non zero result are

$$- \left(e^{\frac{i}{\sqrt{2\alpha'}}\tilde{\phi}} \Big|_{\infty} \partial\phi \Big|_1 e^{-\frac{i}{\sqrt{2\alpha'}}\phi} \Big|_0 \right) - \left(e^{-\frac{i}{\sqrt{2\alpha'}}\tilde{\phi}} \Big|_{\infty} \partial\phi \Big|_1 e^{\frac{i}{\sqrt{2\alpha'}}\phi} \Big|_0 \right) , \quad (6.8)$$

so that Eq. (6.7) reduces to

$$\begin{aligned} T_{\phi} \sim & - \left\langle e^{\frac{i}{\sqrt{2\alpha'}}\tilde{\phi}} \Big|_{\infty} \partial\phi \Big|_1 e^{-\frac{i}{\sqrt{2\alpha'}}\phi} \Big|_0 \text{Tr} \left(\sigma^1 e^{\frac{iA}{2\sqrt{2\alpha'}}} \oint d\sigma \partial_{\sigma} \phi \otimes \sigma^1 \right) \right\rangle \\ & - \left\langle e^{-\frac{i}{\sqrt{2\alpha'}}\tilde{\phi}} \Big|_{\infty} \partial\phi \Big|_1 e^{\frac{i}{\sqrt{2\alpha'}}\phi} \Big|_0 \text{Tr} \left(\sigma^1 e^{\frac{iA}{2\sqrt{2\alpha'}}} \oint d\sigma \partial_{\sigma} \phi \otimes \sigma^1 \right) \right\rangle . \end{aligned} \quad (6.9)$$

To proceed further, we first notice that $\partial\phi$ basically reads the momentum carried by the second exponential factor inside the correlators of Eq. (6.9); hence T_{ϕ} becomes

$$T_{\phi} \sim \left\langle \left[e^{-\frac{i}{\sqrt{2\alpha'}}(q-\tilde{q})} - e^{\frac{i}{\sqrt{2\alpha'}}(q-\tilde{q})} \right] \text{Tr} \left(\sigma^1 e^{\frac{iA}{2\sqrt{2\alpha'}}} \oint d\sigma \partial_{\sigma} \phi \otimes \sigma^1 \right) \right\rangle , \quad (6.10)$$

where q and $\tilde{q}$ are the zero modes of ϕ and $\tilde{\phi}$. Let us recall that in our conventions ²¹, a vertex operator describing a closed string state with Kaluza-Klein index n_{ϕ} and winding number w_{ϕ} , contains the term

$$e^{ik_L\phi} e^{k_R\tilde{\phi}} \sim e^{\frac{i}{\sqrt{2\alpha'}}(n_{\phi} + \frac{w_{\phi}}{2})q_{\phi}} e^{\frac{i}{\sqrt{2\alpha'}}(n_{\phi} - \frac{w_{\phi}}{2})\tilde{q}_{\phi}} \quad (6.11)$$

According to this formula, we easily see that the two terms in the square brackets of Eq. (6.10) correspond respectively to $(n_{\phi} = 0, w_{\phi} = 2)$ and $(n_{\phi} = 0, w_{\phi} = -2)$. Therefore, even if the left-right identification induced by the disk topology of the amplitude implies that both terms are simply equal to one and give no direct contribution, their presence is not trivial. In fact, since they carry different winding numbers, they give a different contribution to the contour integral $\oint \partial\phi$ because

$$\frac{A}{2\sqrt{2\alpha'}} \oint d\sigma \partial_{\sigma} \phi = \pi \frac{w_{\phi}}{2} A \quad (6.12)$$

²¹Our conventions on closed string fields are slightly different from the one of Ref. [13]; in particular the left and the right components of the coordinate X^{μ} are defined as $X^{\mu}(z, \bar{z}) = \frac{1}{2}[X^{\mu}(z) + X^{\mu}(\bar{z})]$, where $z = e^{2(\tau+i\sigma)}$. This implies that k_L and k_R momenta are quantized as follows

$$k_L^{\mu} = \frac{1}{2} \left(\frac{n^{\mu}}{R} + \frac{w^{\mu}R}{\alpha'} \right) , \quad k_R^{\mu} = \frac{1}{2} \left(\frac{n^{\mu}}{R} - \frac{w^{\mu}R}{\alpha'} \right) .$$

At this point the evaluation of Eq. (6.10) simply reduces to that of the trace factor and one gets

$$T_\phi \sim \left[-\text{Tr}(\sigma^1 e^{iA\pi\sigma^1}) + \text{Tr}(\sigma^1 e^{-iA\pi\sigma^1}) \right] \sim \sin(\pi A) ; \quad (6.13)$$

this result is zero for $A = 1$ showing that, for this particular background, the “tachyon” tadpole is still vanishing, even if the radius has been deformed away from the critical value.

This same approach can be used to study how the mass² of the “tachyonic” state (5.10) is modified by a radius deformation; in this case, the relevant amplitude is the two-point function in presence of an operator $\mathcal{V}_R$, namely

$$m_\phi = \langle T(\mathcal{V}_T(x) \mathcal{V}_T(y) \mathcal{V}_R(z)) \rangle_A \quad (6.14)$$

Exploiting the projective invariance to fix $y = 1$ and $z = i$, performing the change of variable of Eq. (6.6) and using the form of the vertex operators in terms of ϕ and $\tilde{\phi}$, we obtain

$$m_\phi \sim \langle \left(e^{\frac{i}{\sqrt{2\alpha'}}\tilde{\phi}} - e^{-\frac{i}{\sqrt{2\alpha'}}\tilde{\phi}} \right) \Big|_\infty \partial\phi \Big|_1 \partial\phi \Big|_x \left(e^{\frac{i}{\sqrt{2\alpha'}}\phi} - e^{-\frac{i}{\sqrt{2\alpha'}}\phi} \right) \Big|_0 \text{Tr} \left(e^{\frac{iA}{2\sqrt{2\alpha'}}} \oint d\sigma \partial_\sigma \phi \otimes \sigma^1 \right) \rangle , \quad (6.15)$$

where the Chan-Paton factors of the two open strings cancel out, since $(\sigma^1)^2 = 1$. As before, some terms in this equation are killed by ϕ -momentum conservation and one is left with only two terms

$$m_\phi \sim \langle \left(\left[e^{-\frac{i}{\sqrt{2\alpha'}}(q_\phi - \tilde{q}_\phi)} \right] + \left[e^{\frac{i}{\sqrt{2\alpha'}}(q_\phi - \tilde{q}_\phi)} \right] \right) \text{Tr} \left(e^{\frac{iA}{2\sqrt{2\alpha'}}} \oint d\sigma \partial_\sigma \phi \otimes \sigma^1 \right) \rangle . \quad (6.16)$$

Note that the sign between the two exponential factors is different from that of Eq. (6.10), because now we have two $\partial\phi$ insertions. Evaluating the trace over the Chan-Paton factors, one finally obtains

$$m_\phi \sim \cos(\pi A) . \quad (6.17)$$

This simple result contains a crucial information: the mass behavior of our “tachyonic” state is very different if one deforms the compactification radius around the trivial configuration $A = 0$ or around the kink solution $A = 1$; indeed

$$\langle T(\mathcal{V}_T(x) \mathcal{V}_T(y) \mathcal{V}_R(z)) \rangle_{A=0} = -\langle T(\mathcal{V}_T(x) \mathcal{V}_T(y) \mathcal{V}_R(z)) \rangle_{A=1} . \quad (6.18)$$

We already know that in the absence of a $\mathcal{T}$ background, the D1/anti-D1 pair develops a tachyonic instability when we take the decompactification limit $R \rightarrow \infty$;

this fact can now be seen directly using the result we have just obtained. Indeed, from Eq. (5.2), it is easy to realize that the mass² of the two lightest modes ($n = 0$, $n = -1$) considered in the previous section, *decreases* when the radius R is increased and becomes negative above the critical value R_c . On the other hand, Eq. (6.18) tells us that this behavior is reversed when $A = 1$ and that the system is stable if $R \geq R_c$. Thus, for $A = 1$ one can take the decompactification limit without encountering any tachyonic instability.

Even if here we have considered only the effects of a first order variation of the radius (that is only one $\mathcal{V}_R$ was inserted), the results we have obtained are quite general and a similar pattern is found also when several $\mathcal{V}_R$ insertions are taken into account. In fact, as is clear from the two explicit examples just described, the relevant part in the disk amplitudes describing the effect of a radius deformation comes from the factor

$$e^{\frac{iA}{2\sqrt{2\alpha'}} \oint d\sigma \partial_\sigma \phi \otimes \sigma^1} = \cos\left(\pi \frac{w_\phi}{2} A\right) + \sigma^1 \sin\left(\pi \frac{w_\phi}{2} A\right), \quad (6.19)$$

which has to be inserted in the correlation functions according to Eq. (6.5). In particular, $\mathcal{V}_R$ can only give rise to combination with even winding number (just like in Eq. (6.16)), so that the $\sin(\dots)$ term always vanishes with our preferred value $A = 1$. On the other hand, the $\cos(\dots)$ term simply changes the sign in front of operators with odd $w_\phi/2$. It is very instructive to use this input directly in the vertex describing the marginal deformation of the radius. To this aim, we can focus on Eq. (6.2), since, after the insertion of the first $\mathcal{V}_R$ in the $(-1, -1)$ picture, the superghost anomaly of the disk is compensated and for all other insertions one should use vertex $\mathcal{V}_R$ in the $(0, 0)$. In the (ϕ, η) language this vertex reads

$$\begin{aligned} \mathcal{V}_R &= \partial X(z) \bar{\partial} \tilde{X}(\bar{z}) = -(2\alpha') \xi \eta \tilde{\xi} \tilde{\eta} \\ &= \sqrt{2\alpha'} \eta \tilde{\eta} \left[e^{\frac{i}{\sqrt{2\alpha'}} \phi} + e^{-\frac{i}{\sqrt{2\alpha'}} \phi} \right] \left[e^{\frac{i}{\sqrt{2\alpha'}} \tilde{\phi}} + e^{-\frac{i}{\sqrt{2\alpha'}} \tilde{\phi}} \right] \\ &= \sqrt{2\alpha'} \eta \tilde{\eta} \left(e^{\frac{i}{\sqrt{2\alpha'}} (\phi - \tilde{\phi})} + e^{-\frac{i}{\sqrt{2\alpha'}} (\phi - \tilde{\phi})} + e^{\frac{i}{\sqrt{2\alpha'}} (\phi + \tilde{\phi})} + e^{-\frac{i}{\sqrt{2\alpha'}} (\phi + \tilde{\phi})} \right). \end{aligned} \quad (6.20)$$

Since the Wilson line $A = 1$ simply changes the sign in front of the exponentials with $\pm(\phi - \tilde{\phi})$, we can forget the presence of the non-trivial background in the trace factor, provided that we use the following modified vertex operator $\mathcal{V}'_R$

$$\begin{aligned} \mathcal{V}'_R &= \sqrt{2\alpha'} \eta \tilde{\eta} \left(e^{\frac{i}{\sqrt{2\alpha'}} (\phi - \tilde{\phi})} - e^{-\frac{i}{\sqrt{2\alpha'}} (\phi - \tilde{\phi})} + e^{\frac{i}{\sqrt{2\alpha'}} (\phi + \tilde{\phi})} - e^{-\frac{i}{\sqrt{2\alpha'}} (\phi + \tilde{\phi})} \right) \\ &= \sqrt{2\alpha'} \eta \tilde{\eta} \left[e^{\frac{i}{\sqrt{2\alpha'}} \phi} - e^{-\frac{i}{\sqrt{2\alpha'}} \phi} \right] \left[e^{\frac{i}{\sqrt{2\alpha'}} \tilde{\phi}} - e^{-\frac{i}{\sqrt{2\alpha'}} \tilde{\phi}} \right]. \end{aligned} \quad (6.21)$$

If one uses as independent variables the fields ϕ' and ξ given in Eq. (5.5), this expression takes a very simple form

$$\mathcal{V}'_R = -\left(i\sqrt{2\alpha'} \eta \psi\right)\left(i\sqrt{2\alpha'} \tilde{\eta} \tilde{\psi}\right) = -\partial\phi' \bar{\partial}\tilde{\phi}' . \quad (6.22)$$

Thus, the presence of the “tachyon” vacuum expectation value $A = 1$ effectively substitutes X^1 with ϕ' in the definition of the marginal operator related to radius deformations and transforms it into $-\partial\phi' \bar{\partial}\tilde{\phi}'$. The minus sign in this expression has a crucial importance; in fact, this means that increasing the radius of the X^1 coordinate ($R_c \rightarrow R_c + \delta R$) is equivalent, at $A = 1$, to *decrease* the radius of the ϕ' coordinate ($R_c \rightarrow R_c - \delta R$), so that the decompactification limit can be also seen as a $R_{\phi'} \rightarrow 0$ limit. It is by now well known that the behavior of open strings compactified on a small circle is more easily understood by making a T-duality transformation. In our case, the field ϕ' satisfies Neumann boundary conditions like ϕ , since both come from the bosonization procedure described in Appendix A; therefore after a T-duality, ϕ' becomes a Dirichlet coordinate ϕ'_D compactified on a circle of radius $R_{\phi'_D} = \alpha'/R_\phi$ and the decompactification limit of the original space-time translates into the $R_{\phi'_D} \rightarrow \infty$ limit. Thus in Minkowski space the bound state between a D1 brane and its anti-brane is described by one Neumann field (X^0) and by nine Dirichlet coordinates ($\phi'_D, X^2, \dots X^9$). This is an indication that, even if the starting point of our analysis was a system with only a $SO(8)$ symmetry manifest, the stable configuration in the flat space has an enlarged $SO(9)$ symmetry; in fact, even if we do not show it explicitly here, it is not difficult to see that ϕ' stays on the same footing of other Dirichlet coordinates: for instance, the full spectrum of open strings living on the bound state is invariant under the exchange $(\phi'_D, \xi) \leftrightarrow (X^2, \psi^2)$. The $SO(9)$ symmetry of the final configuration in the decompactified space corresponds to the little group of massive particle living in ten dimensions, so that the resulting configuration can be interpreted as a D-particle in Type I theory.

Since now we know the effect of radius deformation on the conformal theory living on the D1/anti-D1 system after tachyon condensation, we can return to the problem of the mass M of the new D-particle state and show that for $A = 1$ it does not depend on the radius. This can be shown by means of the same approach used by J. Polchinski in Ref. [4] to find the mass of the D-branes in Type II theories. One starts by computing the vacuum energy between two identical systems and then extracts the part corresponding to the exchange of closed strings in the NS-NS spin structure. In the limit of very large separation, only the massless states

contribute and this string result, which depends only on α' and g , has to be equal to the field theory answer which instead is given in terms of the mass M and of the Newton's constant G . Using this method, we now check that, in the absence of a $\mathcal{T}$ background ($A = 0$), the interaction between two D1/anti-D1 pairs give the expected answer. Newton's law (in ten dimension) tells that the potential energy between two masses M placed at a distance ℓ from each other scales as

$$U \sim G \frac{M^2}{\ell^6}, \quad (6.23)$$

where the Newton's constant is proportional to $1/R$ since one dimension (X^1) is compactified. Also the string result scales as $1/\ell^6$; in fact, the vacuum energy between two p -extended objects at a distance ℓ from each other is

$$U_{\text{string}}^{A=0} \sim V_{p+1} \frac{1}{\ell^{7-p}} \quad (6.24)$$

where V_{p+1} is the world-volume of the p -brane. In our case $p = 1$ and V_2 scales as R . Thus, by equating the R dependence of Eqs. (6.23) and (6.24) for $p = 1$, we find, as expected, that the mass of the system is proportional to the radius: $M \sim R$. In the presence of the kink solution ($A = 1$), the gravitational interaction is still described by Eq. (6.23), but the string computation is different. In fact, from Eq. (6.22) we know that the effective radius R'_ϕ one should use in the conformal field theory analysis is the inverse of the physical radius R along the X^1 direction. Thus, for $p = 1$ we get

$$U_{\text{string}}^{A=1} \sim V'_2 \frac{1}{\ell^6} \sim \frac{1}{R} \frac{1}{\ell^6} \quad (6.25)$$

Equating the R behavior of Eqs. (6.23) and (6.25) we now find that at $A = 1$ the mass M of the D1/ant-D1 pair is *independent* of R . Thus, the mass of the Type I D-particle described by this system in the flat Minkowski space is the same as the one computed at the critical radius, *i.e.*

$$M = \frac{1}{\sqrt{\alpha'} g_I} \quad (6.26)$$

as we have seen in Eq. (5.16).

In these notes we have mainly concentrated on the construction of the stable non-BPS D-particle of Type I which is predicted and required by the heterotic/Type I duality. However, the techniques we have used and the strategy we have discussed can be generalized and applied to other systems and theories. Even if the entire construction seems quite complicated and technically involved, the emerging picture

is very satisfactory and appealing and makes manifest the fact that the presence of a tachyonic field in a string theory is not necessarily a sign of inconsistency. Of course there are many issues and developments that have not been covered in these notes but which are very interesting, such as for example the construction of other stable non-BPS D-branes of Type I [25, 40] or the relation between the D-brane charge and K-theory [25, 26]. Moreover, there are still some open problems which remain to be solved: among them we can mention the problem of understanding the mechanism of tachyon condensation directly at the level of the effective theory of the branes which in turn requires also a better understanding of the fate of the various gauge fields living on D-brane world volume [25, 44]. A progress in this direction presumably could help to understand the mechanism of tachyon condensation also in those cases where the tachyon appears in the closed string spectrum.

Acknowledgements

It is a pleasure to thank P. Di Vecchia, M. Frau, L. Gallot, I. Pesando, S. Sciuto and P. Strigazzi for many useful discussions, and the organizers and the participants of the School *Quantum Aspects of Gauge Theories, Supersymmetry and Unification* in Leuven (Belgium) January 1999 for their stimulating questions and observations. R. Russo wishes also to thank A. Bilal and D. Matalliotakis for many valuable suggestions and observations.

Appendix A

In Section 5 we have extensively used the well known equivalence between a compact bosonic field and two real fermions. This equivalence is exploited in string theory in many contexts, but, even if the basic idea is always encoded by Eq. (5.3), there are some differences among various cases. For instance, the bosonization procedure relates the two formulations of heterotic theory mentioned in Section 2; however, in that case, the bosonic fields satisfied periodic boundary conditions and the critical radius was $\sqrt{\alpha'}$. On the other hand, if Neumann boundary conditions are imposed, the equivalence holds for $R = \sqrt{2\alpha'}$. In fact, by looking at Eq. (5.3), it is clear that

the first fermionic state one can construct by acting with $(\xi \pm i\eta)$ on the vacuum corresponds, in the bosonic description, to a scalar state of momentum $\pm 1/\sqrt{2\alpha'}$. Since higher fermionic states are obtained by introducing other factors of $(\xi \pm i\eta)$, in the bosonic language one has to consider states whose momentum k is $n/\sqrt{2\alpha'}$ with $n \in \mathbb{Z}$. Further evidence that the two formulations really describe the same theory is provided by the computation of the partition function. In terms of the field X^1 , the partition function factorizes in a sum over the allowed momenta and in the oscillator contribution

$$\text{Tr} \left(q^{2L_0} \right) = \sum q^{2\alpha' k^2} \prod_{n=1}^{\infty} \frac{1}{(1 - q^{2n})} . \quad (\text{A.1})$$

However, because of the particular form of the momentum, the sum take a very simple form and can be rewritten as a product

$$\sum_{n \in \mathbb{Z}} q^{n^2} = \prod_{n=1}^{\infty} (1 - q^{2n}) (1 + q^{2n-1})^2 , \quad (\text{A.2})$$

and so Eq. (A.1) can be recast in a different form

$$\sum q^{2\alpha' k^2} \prod_{n=1}^{\infty} \frac{1}{(1 - q^{2n})} = \prod_{n=1}^{\infty} (1 + q^{2n-1})^2 , \quad (\text{A.3})$$

that directly matches the partition function of two real NS fermions. However, this is not exactly the result we needed in Section 5, where the bosonic field was compactified on a circle of radius $R_c = \sqrt{\alpha'}/2$ and could have integer or half-integer values of the momentum according to various Chan-Paton factors (but, of course not both together). If one looks at the states proportional to σ^1 or σ^2 , it is easy to realize that they simply are a subset of those that can be constructed when $R = \sqrt{2\alpha'}$; in particular their momentum is of the form $(2n + 1)/\sqrt{2\alpha'}$. This consideration suggests that, at the critical radius we are interested in, the only bosonic states that can be constructed correspond to fermionic states with an *odd* number of ξ and η oscillators. On the other hand, the states proportional to σ^3 or $\mathbb{1}$ are periodic: this implies that their momentum is integer in terms of R_c and thus it can take only even values in terms of $R = \sqrt{2\alpha'}$. In this case the corresponding fermionic states contain an *even* number of ξ and η oscillators. Thus it turns out that the states with half-integer momentum of the bosonic field correspond to states with the usual GSO projection in the fermionic language; on the contrary, when the momentum is quantized in integer multiples of $1/R_c$, one reproduces the sector with the "wrong" GSO projection of the two real fermions. This can be checked again

at the level of partition function. Let us focus, for example, on the $\sigma^1 \sigma^2$ sector. Here the sum over the allowed momenta in Eq. (A.1) takes a slightly different form, but nevertheless it can be transformed in the expected partition function by using again the identity (A.2)

$$\begin{aligned} \sum_{n \in \mathbb{Z}} q^{(2n+1)^2} &= \frac{1}{2} \sum_{n \in \mathbb{Z}} (q^{n^2} - (-q)^{n^2}) \\ &= \frac{1}{2} \prod_{n=1}^{\infty} (1 - q^{2n}) \left[(1 + q^{2n-1})^2 - (1 - q^{2n-1})^2 \right]. \end{aligned} \quad (\text{A.4})$$

It is possible to resume this result in a simple requirement, which also allows us to treat the states with different Chan-Paton factors in the same way. In fact, we can think that the momentum of the bosonic field X^1 is always allowed to assume both integer and half-integer values, but, then, we introduce a new discrete symmetry h that selects physical states. From previous analysis, it is clear that h acts on ξ and η as a GSO projection; then from Eqs. (5.4)-(5.5) one can derive how h translates in terms of the rebosonized fields. For instance, by taking ϕ and η as independent fields, it is easy to see from Eq. (5.4) that h transforms $: e^{\frac{i}{\sqrt{2\alpha'}} \phi} :$ in $(- : e^{-\frac{i}{\sqrt{2\alpha'}} \phi} :)$. The explicit form of h in the different languages is reported in Table 1.

h				
$\left\{ \begin{array}{l} X \rightarrow X + 2\pi\sqrt{\frac{\alpha'}{2}} \\ \psi^\mu \rightarrow \psi^\mu \\ 0\rangle_{-1} \rightarrow 0\rangle_{-1} \\ \lambda \rightarrow \sigma^3 \lambda \sigma^3 \end{array} \right\}$	$\Leftrightarrow$	$\left\{ \begin{array}{l} \xi \rightarrow -\xi \\ \eta \rightarrow -\eta \\ \psi^\mu \rightarrow \psi^\mu \\ 0\rangle_{-1} \rightarrow 0\rangle_{-1} \\ \lambda \rightarrow \sigma^3 \lambda \sigma^3 \end{array} \right\}$	$\Leftrightarrow$	$\left\{ \begin{array}{l} \phi \rightarrow -\phi + 2\pi\sqrt{\frac{\alpha'}{2}} \\ \eta \rightarrow -\eta \\ \psi^{\mu \neq 1} \rightarrow \psi^{\mu \neq 1} \\ 0\rangle_{-1} \rightarrow 0\rangle_{-1} \\ \lambda \rightarrow \sigma^3 \lambda \sigma^3 \end{array} \right\}$

Table 1: The discrete symmetry h that defines the physical states according to the different Chan-Paton factors: physical states are invariant under these transformations.

In a similar way one can also write down the GSO projection in terms of sets of independent fields and derive the results of Table 2.

Of course, the boundary conditions of the new bosonic field ϕ depends on those of ξ and ψ . For instance, if both fermions are in the NS sector, it is sufficient to read Eq. (A.3) backwards to see that ϕ is an usual Neumann field. However, while ξ is

$(-1)^F$				
$\left\{ \begin{array}{l} X \rightarrow X \\ \psi^\mu \rightarrow -\psi^\mu \\ 0\rangle_{-1} \rightarrow - 0\rangle_{-1} \\ \lambda \rightarrow \sigma^3 \lambda \sigma^3 \end{array} \right\}$	$\Leftrightarrow$	$\left\{ \begin{array}{l} \xi \rightarrow \xi \\ \eta \rightarrow \eta \\ \psi^\mu \rightarrow -\psi^\mu \\ 0\rangle_{-1} \rightarrow - 0\rangle_{-1} \\ \lambda \rightarrow \sigma^3 \lambda \sigma^3 \end{array} \right\}$	$\Leftrightarrow$	$\left\{ \begin{array}{l} \phi \rightarrow -\phi \\ \eta \rightarrow \eta \\ \psi^{\mu \neq 1} \rightarrow -\psi^{\mu \neq 1} \\ 0\rangle_{-1} \rightarrow - 0\rangle_{-1} \\ \lambda \rightarrow \sigma^3 \lambda \sigma^3 \end{array} \right\}$

Table 2: The generalized GSO projection defined in Eq. (4.9) written in terms of different fields: physical states are invariant under these transformations.

always a NS field, ψ can also be in the R sector: in this case the combination of the two gives a bosonic field with Dirichlet boundary conditions on one endpoint and Neumann boundary conditions on the other. Thus, ϕ is half-integer moded and can not carry momentum, thus its partition function is simply $(\prod(1 - q^{2m-1}))^{-1}$. The partition function in the fermionic description can be recast in this form showing again that the two representations are equivalent

$$Z_R = \prod_{m=1}^{\infty} (1 + q^{2m-1}) (1 + q^{2m}) = \prod_{n=1}^{\infty} (1 + q^n) = \prod_{m=1}^{\infty} \frac{1}{(1 - q^{2m-1})}. \quad (\text{A.5})$$

Finally a useful observation is that the two discrete symmetries $(-1)^F$ and h can be combined so that the resulting transformation is much simpler in the (ϕ, η) description.

$(-1)^F h$
$\phi \rightarrow \phi + 2\pi\sqrt{\frac{\alpha'}{2}}$ $\eta \rightarrow -\eta$ $\psi^{\mu \neq 1} \rightarrow -\psi^{\mu \neq 1}$ $ 0\rangle_{-1} \rightarrow - 0\rangle_{-1}$ $\lambda \rightarrow \lambda$

Table 3: The $(-1)^F h$ in terms of ϕ and η

Since the physical states have to be even under both $(-1)^F$ and h separately, they are also even under $(-1)^F h$ and either $(-1)^F$ or h , and indeed the two requirements are equivalent.

References

- [1] M. B. Green, J. H. Schwarz, and E. Witten, *Superstring Theory*, Cambridge University Press, 1987.
- [2] J. H. Schwarz, *An $SL(2, Z)$ Multiplet of Type IIB Superstrings*, Phys. Lett. **B360**, 13 (1995), [hep-th/9508143](#).
- [3] J. Polchinski and E. Witten, *Evidence for Heterotic-Type I string duality*, Nucl. Phys. **B460**, 525 (1996), [hep-th/9510169](#).
- [4] J. Polchinski, *Dirichlet branes and Ramond-Ramond charges*, Phys. Rev. Lett. **75**, 184 (1995), [hep-th/9510017](#).
- [5] J. Polchinski, “*TASI lectures on D-branes*”, lectures given at the TASI summer school (1996) [hep-th/9611050](#).
- [6] A. Sen, *$SO(32)$ Spinors of Type I and Other Solitons on Brane-Antibrane Pair*, JHEP **9809**, 023 (1998), [hep-th/9808141](#).
- [7] A. Sen, *Type I D-particle and its Interactions*, JHEP **9810**, 021 (1998), [hep-th/9809111](#).
- [8] A. Dabholkar, *Ten Dimensional Heterotic String as a Soliton*, Phys. Lett **B357**, 307 (1995), [hep-th/9506160](#).
- [9] A. A. Tseytlin, *Heterotic-type I superstring duality and low-energy effective actions*, Nucl. Phys. **B467**, 383 (1995), [hep-th/9512081](#).
- [10] D. Matalliotakis, H. P. Nilles, and S. Theisen, *Matching the BPS Spectra of Heterotic - Type I - Type I'*, Phys. Lett. **B421**, 169 (1998), [hep-th/9710247](#).
O. Bergman, M.R. Gaberdiel, and G. Lifschytz, *String Creation and Heterotic-Type I' Duality*, Nucl. Phys. **B524**, 524 (1998), [hep-th/9711098](#).
- [11] C. Bachas, C. Fabre, E. Kiritsis, N. Obers, and P. Vanhove, *Heterotic/type I-duality and D-brane instantons*, Nucl. Phys. **B509**, 33 (1998), [hep-th/9707126](#).
C. Bachas and E. Kiritsis, *F^4 Terms in $N=4$ String Vacua*, Nucl. Phys. [Proc. Suppl.] **55B**, 194 (1997), [hep-th/9611205](#).
E. Kiritsis and N. Obers, *Heterotic/Type-I Duality in $D < 10$ Dimensions, Threshold Corrections and D-Instantons*, JHEP **10**, 004 (1997), [hep-th/9709058](#).

- [12] A. Sen, *Unification of String Dualities*, Nucl. Phys. [Proc. Suppl] **58**, 5 (1997), [hep-th/9609176](#).
- [13] J. Polchinski, *String Theory*, Cambridge University Press, 1998.
- [14] L. Susskind T. Banks, *Brane - Anti-Brane Forces*, [hep-th/9511194](#), 1995.
- [15] A. Sen, *Stable Non-BPS States in String Theory*, JHEP **9808**, 007 (1998), [hep-th/9803194](#).
- [16] A. Sen, *Tachyon Condensation on the Brane Antibrane System*, JHEP **9808**, 012 (1998), [hep-th/9805170](#).
- [17] A. Sen, *Stable Non-BPS Bound States of BPS D-branes*, JHEP **9808**, 010 (1998), [hep-th/9805019](#).
- [18] A. Sen, *BPS D-branes on Non-supersymmetric Cycles*, JHEP **9812**, 021 (1998), [hep-th/9812031](#).
- [19] A. Sagnotti, *Non-Perturbative Quantum Field Theory*, Eds G. Mack et al. (Pergamon Press, 1988) 521.
 N. Ishibashi and T. Onogi, *Conformal field theories on surfaces with Boundaries and Crosscaps*, Mod. Phys. Lett. **A4**, 161 (1989).
 G. Pradisi and A. Sagnotti, *Open string Orbifolds* Phys. Lett. **B216**, 59 (1989).
 P. Hořava, *Strings on world sheet Orbifolds* Nucl. Phys. **B327**, 461 (1989).
- [20] O. Bergman and M.R. Gaberdiel, *Stable non-BPS D-particles*, Phys. Lett. **B441**, 133 (1998), [hep-th/9806155](#).
- [21] C. G. Callan, C. Lovelace, C. R. Nappi, and S. A. Yost, *Loop corrections to superstring equations of motion*, Nucl. Phys. **B308**, 221 (1988).
- [22] P. Di Vecchia, M. Frau, I. Pesando, S. Sciuto, A. Lerda, and R. Russo, *Classical p-branes from boundary state*, Nucl. Phys. **B507**, 259 (1997), [hep-th/9707068](#).
 M. Billò, P. Di Vecchia, M. Frau, A. Lerda, I. Pesando, R. Russo, and S. Sciuto, *Microscopic string analysis of the D0-D8 brane system and dual R-R states*, Nucl. Phys. **B526**, 1998 (1998), [hep-th/9802088](#).
 M. Billò, P. Di Vecchia, M. Frau, A. Lerda, R. Russo, and S. Sciuto, *The Lorentz Force between D0 and D6 Branes in String and M(atrrix) Theory*, Mod. Phys. Lett. **A13**, 2977 (1998), [hep-th/9805091](#).

- [23] O. Bergman and M.R. Gaberdiel, *Non-BPS States in Heterotic-Type IIA Duality*, JHEP **9903**, 013 (1999), [hep-th/9901014](#).
- [24] A. Sen, *Non-BPS States and Branes in String Theory*, [hep-th/9904207](#), 1999.
- [25] E. Witten, *D-Branes and K-Theory*, JHEP **9812**, 019 (1998), [hep-th/9503124](#).
- [26] P. Hořava, *Type IIA D-Branes, K-Theory, and Matrix Theory*, [hep-th/9812135](#), 1998.
- [27] K. Dasgupta and J. Park, *Brane mapping under $(-1)^{F_L}$* , [hep-th/9903125](#), 1999.
- [28] H. Garcia-Compeán, *D-branes in Orbifold Singularities and Equivariant K-Theory*, [hep-th/9812226](#), 1998.
- [29] S. Gukov, *K-Theory, Reality, and Orientifolds*, [hep-th/9901042](#), 1999.
- [30] K. Olsen and R.J. Szabo, *Brane Descent Relations in K-theory*, [hep-th/9904153](#), 1999.
- [31] D. Friedan, D. Martinec, and S. Shenker, *Conformal invariance, supersymmetry and string theory*, Nucl. Phys. **B271**, 93 (1986).
- [32] A. Dabholkar and J. Harvey, *Nonrenormalization of the superstring tension*, Phys. Rev. Lett. **63**, 478 (1989).
- [33] M. B. Green and J. H. Schwarz, *Anomaly cancellation in supersymmetric $D = 10$ gauge theory and superstring theory*, Phys. Lett. **149B**, 117 (1984).
- [34] J. Polchinski and Y. Cai, *Consistency of open superstring theories*, Nucl. Phys. **B286**, 91 (1988).
- [35] C. G. Callan, C. Lovelace, C. R. Nappi, and S. A. Yost, *Adding holes and crosscaps to the superstring*, Nucl. Phys. **B293**, 83 (1987).
- [36] E. Witten, *String theory dynamics in various dimension*, Nucl. Phys. **B443**, 85 (1995), [hep-th/9503124](#).
- [37] I. R. Klebanov and A. A. Tseytlin, *D-Branes and Dual Gauge Theories in Type 0 String Theory*, Nucl. Phys. **B547**, 143 (1999), [hep-th/9811035](#).

- [38] I. Pesando, *On the Effective Potential of the Dp - anti- Dp system in type II theories*, [hep-th/9902181](#), 1999.
- [39] J. Polchinski and A. Strominger, *New Vacua for Type II String Theory*, Phys. Lett. **B388**, 736 (1996), [hep-th/9510227](#).
E. Gava, K. Narain and M. Sarmadi *On the Bound States of p- and (p+2)-Branes*, Nucl. Phys. **B504**, 214 (1997), [hep-th/9704006](#).
- [40] M. Frau, L. Gallot, A. Lerda, and P. Strigazzi, *Stable non-BPS D-branes in Type I string theory*, [hep-th/9903123](#), 1999.
- [41] E. G. Gimon and J. Polchinski, *Consistency Conditions for Orientifolds and D-Manifolds*, Phys. Rev. **D54**, 1667 (1996), [hep-th/9601038](#).
- [42] M. B. Green, *Space-time duality and Dirichlet string theory*, Phys. Lett. **B266**, 325 (1991).
- [43] M. Ademollo, A. D’Adda, R. D’Auria, E. Napolitano, P. Di Vecchia, F. Gliozzi, and S. Sciuto, *Unified dual model for interacting open and closed strings*, Nucl. Phys. **B77**, 189 (1974).
- [44] P. Yi, *Membranes from Five-Branes and Fundamental Strings from Dp Branes*, [hep-th/9901159](#), 1999.